

# A SURVEY OF THE SPECTRAL AND DIFFERENTIAL GEOMETRIC ASPECTS OF THE DE RHAM-HODGE-SKRYPNIK THEORY RELATED WITH DELSARTE TRANSMUTATION OPERATORS IN MULTIDIMENSION AND ITS APPLICATIONS TO SPECTRAL AND SOLITON PROBLEMS. PART 2

Y.A. PRYKARPATSKY<sup>\*)\*\*</sup>), A.M. SAMOILENKO<sup>\*\*</sup>), AND A. K. PRYKARPATSKY<sup>\*\*\*</sup>)

**ABSTRACT.** The differential-geometric and topological structure of Delsarte transmutation operators and associated with them Gelfand-Levitan-Marchenko type equations are studied making use of the De Rham-Hodge-Skrypnik differential complex. The relationships with spectral theory and special Berezansky type congruence properties of Delsarte transmuted operators are stated. Some applications to multidimensional differential operators are done including three-dimensional Laplace operator, two-dimensional classical Dirac operator and its multidimensional affine extension, related with self-dual Yang-Mills equations. The soliton like solutions to the related set of nonlinear dynamical systems are discussed.

## 1. DE RHAM-HODGE-SKRYPNIK THEORY ASPECTS AND RELATED DELSARTE-DARBOUX TYPE BINARY TRANSFORMATIONS

**1.1** A differential-geometric analysis of Delsarte-Darboux type transformations that was done in Chapter 3 of Part 1 for differential operator expressions acting in a functional space  $\mathcal{H} = L_1(T; H)$ , where  $T = \mathbb{R}^2$  and  $H := L_2(\mathbb{R}^2; \mathbb{C}^2)$ , appears to have a deep relationship with classical de Rham-Hodge-Skrypnik theory [3, 4, 5, 6] devised in the midst of the past century for a set of commuting differential operators defined, in general on a smooth compact  $m$ -dimensional metric space  $M$ . Concerning our problem of describing the spectral structure of Delsarte-Darboux type transformations acting in  $\mathcal{H}$ , we preliminarily consider following Part 1 some backgrounds of the de-Rham-Hodge-Skrypnik differential complex theory devised for studying transformations of differential operators. Consider a smooth metric space  $M$  being a suitably compactified form of the space  $\mathbb{R}^m$ ,  $m \in \mathbb{Z}_+$ . Then one can define on  $M_T := T \times M$  the standard Grassmann algebra  $\Lambda(M_T; \mathcal{H})$  of differential forms on  $T \times M$  and consider a generalized external I.V. Skrypnik [3, 4] anti-differentiation operator  $d_{\mathcal{L}} : \Lambda(M_T; \mathcal{H}) \rightarrow \Lambda(M_T; \mathcal{H})$  acting as follows: for

---

1991 *Mathematics Subject Classification.* Primary 34A30, 34B05 Secondary 34B15  
PACS: 02.30.Jr, 02.30.Uu, 02.30.Zz, 02.40.Sf.

*Key words and phrases.* Delsarte transmutation operators, De Rham-Hodge-Skrypnik differential complex, Darboux transformations, Gelfand-Levitan-Marchenko equations, Dirac operator, Laplace operator, soliton like solutions.

The third author was supported in part by a local AGH grant.

any  $\beta^{(k)} \in \Lambda^k(M_T; \mathcal{H})$ ,  $k = \overline{0, m}$ ,

$$(1) \quad d_{\mathcal{L}}\beta^{(k)} := \sum_{j=1}^2 dt_j \wedge L_j(t; x|\partial)\beta^{(k)} \in \Lambda^{k+1}(M_T; \mathcal{H}),$$

where, by definition,

$$(2) \quad L_j(t; x|\partial) := \partial/\partial t_j - L_j(t; x|\partial)$$

$j = \overline{1, 2}$ , are suitably defined linear differential operators in  $\mathcal{H}$ , commuting to each other, that is

$$(3) \quad [L_1, L_2] = 0.$$

We will put, in general, that differential expressions

$$(4) \quad L_j(t; x|\partial) := \sum_{|\alpha|=0}^{n_j(L)} a_{\alpha}^{(j)}(t; x) \frac{\partial^{|\alpha|}}{\partial x^{\alpha}},$$

with coefficients  $a_{\alpha}^{(j)} \in C^1(T; C^{\infty}(M; End\mathbb{C}^N))$ ,  $|\alpha| = \overline{0, n_j(L)}$ ,  $n_j^{\alpha} \in \mathbb{Z}_+$ ,  $j = \overline{0, 1}$ , are some closed normal densely defined operators in the Hilbert space  $H$  for any  $t \in T$ . It is easy to observe that the anti-differentiation of  $d_{\mathcal{L}}$  defined by (1) is a generalization of the usual external anti-differentiation

$$(5) \quad d = \sum_{j=1}^m dx_j \wedge \frac{\partial}{\partial x_j} + \sum_{s=1}^2 dt_s \wedge \frac{\partial}{\partial t_s}$$

for which, evidently, commutation conditions

$$(6) \quad \left[\frac{\partial}{\partial x_j}; \frac{\partial}{\partial x_k}\right] = 0, \quad \left[\frac{\partial}{\partial t_s}; \frac{\partial}{\partial t_l}\right] = 0, \quad \left[\frac{\partial}{\partial x_j}; \frac{\partial}{\partial t_s}\right] = 0$$

hold for all  $j, k = \overline{1, m}$  and  $s, l = \overline{1, 2}$ . If now to substitute within (5)  $\partial/\partial x_j \rightarrow A_j$ ,  $\partial/\partial t_s \rightarrow L_s$ ,  $j = \overline{1, m}$ ,  $s = \overline{1, 2}$ , one gets the anti-differentiation

$$(7) \quad d_{\mathcal{A}} := \sum_{j=1}^m dx_j \wedge A_j(t; x|\partial) + \sum_{s=1}^2 dt_s \wedge L_s(t; x|\partial),$$

where the differential expressions  $A_j, L_s : \mathcal{H} \rightarrow \mathcal{H}$  for all  $j, k = \overline{1, m}$  and  $s, l = \overline{1, 2}$ , satisfy the commutation conditions  $[A_j, A_k] = 0$ ,  $[L_s, L_s] = 0$ ,  $[A_j, L_s] = 0$ , then then operation (7) defines on  $\Lambda(M_T; \mathcal{H})$  an anti-differential with respect to which the co-chain complex.

$$(8) \quad \mathcal{H} \rightarrow \Lambda^0(M_T; \mathcal{H}) \xrightarrow{d_{\mathcal{A}}} \Lambda^1(M_T; \mathcal{H}) \xrightarrow{d_{\mathcal{A}}} \dots \xrightarrow{d_{\mathcal{A}}} \Lambda^{m+2}(M_T; \mathcal{H}) \xrightarrow{d_{\mathcal{A}}} 0$$

is evidently closed, that is  $d_{\mathcal{A}}d_{\mathcal{A}} \equiv 0$ . As the anti-differential (1) is a particular case of (7), we obtain that the corresponding to it co-chain complex (8) is closed too.

**1.2** Below we will follow ideas developed [3, 4, 5, 6, 29]. A differential form  $\beta \in \Lambda(M_T; \mathcal{H})$  will be called  $d_{\mathcal{A}}$ -closed if  $d_{\mathcal{A}}\beta = 0$  and a form  $\gamma \in \Lambda(M_T; \mathcal{H})$  will be called exact or  $d_{\mathcal{A}}$ -homological to zero if there exists on  $M_T$  such a form  $\omega \in \Lambda(M_T; \mathcal{H})$  that  $\gamma = d_{\mathcal{A}}\omega$ .

Consider now the standard [28, 29, 8, 31] algebraic Hodge star-operation

$$(9) \quad * : \Lambda^k(M_T; \mathcal{H}) \rightarrow \Lambda^{m+2-k}(M_T; \mathcal{H}),$$

$k = \overline{0, m+2}$ , as follows: if  $\beta \in \Lambda^k(M_T; \mathcal{H})$ , then the form  $*\beta \in \Lambda^{m+2-k}(M_T; \mathcal{H})$  is such that:

- $(m-k+2)$  - dimensional volume  $|\beta|$  of the form  $*\beta$  equals  $k$ -dimensional volume  $|\beta|$  of the form  $\beta$ ;
- the  $(m+2)$  -dimensional measure  $\bar{\beta}^\tau \wedge *\beta > 0$  under the fixed orientation on  $M_T$ .

Define also on the space  $\Lambda(M_T; \mathcal{H})$  the following natural scalar product: for any  $\beta, \gamma \in \Lambda^k(M_T; \mathcal{H})$ ,  $k = \overline{0, m}$ ,

$$(10) \quad (\beta, \gamma) := \int_{M_T} \bar{\beta}^\tau * \gamma.$$

Subject to the scalar product (10) one can naturally construct the corresponding Hilbert space

$$(11) \quad \mathcal{H}_\Lambda(M_T) := \bigoplus_{k=0}^{m+2} \mathcal{H}_\Lambda^k(M_T)$$

well suitable for our further consideration. Notice also here, that the Hodge star  $*$ -operation satisfies the following easily checkable property: for any  $\beta, \gamma \in \mathcal{H}_\Lambda^k(M_T)$ ,  $k = \overline{0, m}$ ,

$$(12) \quad (\beta, \gamma) = (*\beta, *\gamma),$$

that is the Hodge operation  $*$  :  $\mathcal{H}_\Lambda(M_T) \rightarrow \mathcal{H}_\Lambda(M_T)$  is unitary and its standard adjoint with respect to the scalar product (10) operation satisfies the condition  $(*)' = (*)^{-1}$ .

Denote by  $d'_\mathcal{L}$  the formally adjoint expression to the weak differential operation (1). By means of the operations  $d'_\mathcal{L}$  and  $d_\mathcal{L}$  in the  $\mathcal{H}_\Lambda(M_T)$  one can naturally define [8, 28, 29, 3, 31] the generalized Laplace-Hodge-Skrypnik operator  $\Delta_\mathcal{L} : \mathcal{H}_1(M_T) \rightarrow \mathcal{H}_1(M_T)$  as

$$(13) \quad \Delta_\mathcal{L} = d'_\mathcal{L} d_\mathcal{L} + d'_\mathcal{L} d_\mathcal{L}.$$

Take a form  $\beta \in \mathcal{H}_\Lambda(M_T)$  satisfying the equality

$$(14) \quad \Delta_\mathcal{L} \beta = 0.$$

Such a form is called [3, 29, 31, 8] harmonic. One can also verify that a harmonic form  $\beta \in \mathcal{H}_\Lambda(M_T)$  satisfies simultaneously the following two adjoint conditions:

$$(15) \quad d'_\mathcal{L} \beta = 0, \quad d_\mathcal{L} \beta = 0$$

easily stemming from (13) and (14).

It is easy to check that the following differential operators in  $\mathcal{H}_\Lambda(M_T)$

$$(16) \quad d_\mathcal{L}^* := *d'_\mathcal{L} (*)^{-1}$$

defines also a new external anti-differential operation in  $\mathcal{H}_\Lambda(M_T)$ .

**Lemma 1.** *The corresponding dual to (8) co-chain complex*

$$(17) \quad \mathcal{H} \longrightarrow \Lambda^0(M_T; \mathcal{H}) \xrightarrow{d_\mathcal{L}^*} \Lambda^1(M_T; \mathcal{H}) \xrightarrow{d_\mathcal{L}^*} \dots \xrightarrow{d_\mathcal{L}^*} \Lambda^{m+2}(M_T; \mathcal{H}) \xrightarrow{d_\mathcal{L}^*} 0$$

is exact.

*Proof.* A proof follows owing to the property  $d_\mathcal{L}^* d_\mathcal{L}^* = 0$  holding due to the definition (16).  $\triangleright$

□

**1.3** Denote further by  $\mathcal{H}_{\Lambda(\mathcal{L})}^k(M_T)$ ,  $k = \overline{0, m+2}$ , the cohomology groups of  $d_{\mathcal{L}}$ -closed and by  $\mathcal{H}_{\Lambda(\mathcal{L}^*)}^k(M_T)$ ,  $k = \overline{0, m+2}$ , the cohomology groups of  $d_{\mathcal{L}^*}$ -closed differential forms, respectively, and by  $\mathcal{H}_{\Lambda(\mathcal{L}^*\mathcal{L})}^k(M_T)$ ,  $k = \overline{0, m+2}$ , the abelian groups of harmonic differential forms from the Hilbert sub-spaces  $\mathcal{H}_{\Lambda}^k(M_T)$ ,  $k = \overline{0, m+2}$ . Before formulating next results, define the standard Hilbert-Schmidt rigged chain [12, 13] of positive and negative Hilbert spaces of differential forms

$$(18) \quad \mathcal{H}_{\Lambda,+}^k(M_T) \subset \mathcal{H}_{\Lambda}^k(M_T) \subset \mathcal{H}_{\Lambda,-}^k(M_T),$$

the corresponding hereditary rigged chains of harmonic forms:

$$(19) \quad \mathcal{H}_{\Lambda(\mathcal{L}^*\mathcal{L}),+}^k(M_T) \subset \mathcal{H}_{\Lambda(\mathcal{L}^*\mathcal{L})}^k(M_T) \subset \mathcal{H}_{\Lambda(\mathcal{L}^*\mathcal{L}),-}^k(M_T)$$

and chains of cohomology groups:

$$(20) \quad \begin{aligned} \mathcal{H}_{\Lambda(\mathcal{L}),+}^k(M_T) &\subset \mathcal{H}_{\Lambda(\mathcal{L})}^k(M_T) \subset \mathcal{H}_{\Lambda(\mathcal{L}),-}^k(M_T), \\ \mathcal{H}_{\Lambda(\mathcal{L}^*),+}^k(M_T) &\subset \mathcal{H}_{\Lambda(\mathcal{L}^*)}^k(M_T) \subset \mathcal{H}_{\Lambda(\mathcal{L}^*),-}^k(M_T) \end{aligned}$$

for all  $k = \overline{0, m+2}$ . Assume also that the Laplace-Hodge-Skrypnik operator (13) is reduced upon the space  $\mathcal{H}_{\Lambda}^0(M)$ . Now by reasoning similar to those in [8, 29, 31] one can formulate a little generalized [4, 5, 6, 29] de Rham-Hodge-Skrypnik theorem.

The groups of harmonic forms  $\mathcal{H}_{\Lambda,+}^k(M_T)$ ,  $k = \overline{0, m+2}$ , are, respectively, isomorphic to the homology groups  $(H^k(M_T; \mathbb{C}))^{|\Sigma|}$ ,  $k = \overline{0, m+2}$ , where  $H^k(M_T; \mathbb{C})$  is the  $k$ -th cohomology group of the manifold  $M_T$  with complex coefficients, a set  $\Sigma \subset \mathbb{C}^p$ ,  $p \in \mathbb{Z}_+$ , is the set of suitable "spectral" parameters marking the linear space of independent  $d_{\mathcal{L}^*}$ -closed 0-form from  $\mathcal{H}_{\Lambda(\mathcal{L}),-}^0(M_T)$  and, moreover, the following direct sum decompositions

$$(21) \quad \begin{aligned} \mathcal{H}_{\Lambda,+}^k(M_T) &= \mathcal{H}_{\Lambda(\mathcal{L}^*\mathcal{L}),+}^k(M_T) \oplus \Delta_{\mathcal{L}} \mathcal{H}_{\Lambda,+}^k(M_T) \\ &= \mathcal{H}_{\Lambda(\mathcal{L}^*\mathcal{L}),+}^k(M_T) \oplus d_{\mathcal{L}} \mathcal{H}_{\Lambda,+}^{k-1}(M_T) \oplus d'_{\mathcal{L}} \mathcal{H}_{\Lambda,+}^{k+1}(M_T) \end{aligned}$$

hold for any  $k = \overline{0, m+2}$ .

Another variant of the statement similar to that above was formulated in [3, 4] and reads as the following generalized de Rham-Hodge-Skrypnik theorem.

The generalized cohomology groups  $\mathcal{H}_{\Lambda(\mathcal{L}),+}^k(M_T)$ ,  $k = \overline{0, m+2}$ , are isomorphic, respectively, to the cohomology groups  $(H^k(M_T; \mathbb{C}))^{|\Sigma|}$ ,  $k = \overline{0, m+2}$ .

A proof of this theorem is based on some special sequence [3, 4, 5, 6, 7] of differential Lagrange type identities.▷

Define the following closed subspace

$$(22) \quad \mathcal{H}_0^* := \{\varphi^{(0)}(\eta) \in \mathcal{H}_{\Lambda(\mathcal{L}^*),-}^0(M_T) : d_{\mathcal{L}}^* \varphi^{(0)}(\eta) = 0, \varphi^{(0)}(\eta)|_{\Gamma}, \eta \in \Sigma\}$$

for some smooth  $(m+1)$ -dimensional hypersurface  $\Gamma \subset M_T$  and  $\Sigma \subset (\sigma(L) \cap \bar{\sigma}(L)) \times \Sigma_{\sigma} \subset \mathbb{C}^p$ , where  $\mathcal{H}_{\Lambda(\mathcal{L}^*),-}^0(M_T)$  is, as above, a suitable Hilbert-Schmidt rigged [12, 13] zero-order cohomology group Hilbert space from the co-chain given by (20),  $\sigma(L)$  and  $\sigma(L^*)$  are, respectively, mutual generalized spectra of the sets of differential operators  $L$  and  $L^*$  in  $H$  at  $t = 0 \in T$ . Thereby, the dimension  $\dim \mathcal{H}_0^* = \text{card } \Sigma := |\Sigma|$  is assumed to be known. The next lemma first stated by I.V. Skrypnik [3, 4] is of fundamental meaning for a proof of Theorem 1.2.

**Lemma 2.** *There exists a set of differential  $(k+1)$ -forms  $Z^{(k+1)}[\varphi^{(0)}(\eta), d_{\mathcal{L}}\psi^{(k)}] \in \Lambda^{k+1}(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , and a set of  $k$ -forms  $Z^{(k)}[\varphi^{(0)}(\eta), \psi^{(k)}] \in \Lambda^k(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , parametrized by the set  $\Sigma \ni \eta$ , being semilinear in  $(\varphi^{(0)}(\eta), \psi^{(k)}) \in \mathcal{H}_0^* \times \mathcal{H}_{\Lambda,+}^k(M_T)$ , such that*

$$(23) \quad Z^{(k+1)}[\varphi^{(0)}(\eta), d_{\mathcal{L}}\psi^{(k)}] = dZ^{(k)}[\varphi^{(0)}(\eta), \psi^{(k)}]$$

for all  $k = \overline{0, m+2}$  and  $\eta \in \Sigma$ .

*Proof.* A proof is based on the following Lagrange type identity generalizing that of Part 1 and holding for any pair  $(\varphi^{(0)}(\eta), \psi^{(k)}) \in \mathcal{H}_0^* \times \mathcal{H}_{\Lambda,+}^k(M_T)$ :

$$(24) \quad \begin{aligned} 0 &= \langle d_{\mathcal{L}}^* \varphi^{(0)}(\eta), *(\psi^{(k)} \wedge \bar{\gamma}) \rangle = \langle *d'_{\mathcal{L}}(*)^{-1} \varphi^{(0)}(\eta), *(\psi^{(k)} \wedge \bar{\gamma}) \rangle \\ &= \langle *d'_{\mathcal{L}}(*)^{-1} \varphi^{(0)}(\eta), \psi^{(k)} \wedge \bar{\gamma} \rangle = \\ &= \langle (*)^{-1} \varphi^{(0)}(\eta), d_{\mathcal{L}}\psi^{(k)} \wedge \bar{\gamma} \rangle + Z^{(k+1)}[\varphi^{(0)}(\eta), d_{\mathcal{L}}\psi^{(k)}] \wedge \bar{\gamma} = \\ &= \langle (*)^{-1} \varphi^{(0)}(\eta), d_{\mathcal{L}}\psi^{(k)} \wedge \bar{\gamma} \rangle + dZ^{(k)}[\varphi^{(0)}(\eta), \psi^{(k)}] \wedge \bar{\gamma}, \end{aligned}$$

where  $Z^{(k+1)}[\varphi^{(0)}(\eta), d_{\mathcal{L}}\psi^{(k)}] \in \Lambda^{k+1}(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , and  $Z^{(k)}[\varphi^{(0)}(\eta), \psi^{(k)}] \in \Lambda^k(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , are some semilinear differential forms on  $M_T$  parametrized by a parameter  $\lambda \in \Sigma$ , and  $\bar{\gamma} \in \Lambda^{m+1-k}(M_T; \mathbb{C})$  is arbitrary constant  $(m+1-k)$ -form. Thereby, the semilinear differential  $(k+1)$ -forms  $Z^{(k+1)}[\varphi^{(0)}(\eta), d_{\mathcal{L}}\psi^{(k)}] \in \Lambda^{k+1}(M_T; \mathbb{C})$  and  $k$ -forms  $Z^{(k)}[\varphi^{(0)}(\eta), \psi^{(k)}] \in \Lambda^k(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ ,  $\lambda \in \Sigma$ , constructed above exactly constitute those searched for in the Lemma.  $\square$

**1.4** Based now on Lemma 1.3 one can construct the cohomology group isomorphism claimed in Theorem 1.2 formulated above. Namely, following [3, 4], let us take some singular simplicial [28, 29, 30, 31] complex  $\mathcal{K}(M_T)$  of the compact metric space  $M_T$  and introduce a set of linear mappings  $B_{\lambda}^{(k)} : \mathcal{H}_{\Lambda,+}^k M_T \longrightarrow C_k(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ ,  $\lambda \in \Sigma$ , where  $C_k(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , are free abelian groups over the field  $\mathbb{C}$  generated, respectively, by all  $k$ -chains of singular simplexes  $S^{(k)} \subset M_T$ ,  $k = \overline{0, m+2}$ , from the simplicial complex  $\mathcal{K}(M_T)$ , as follows:

$$(25) \quad B_{\lambda}^{(k)}(\psi^{(k)}) := \sum_{S^{(k)} \in C_k(M_T; \mathbb{C})} S^{(k)} \int_{S^{(k)}} Z^{(k)}[\varphi^{(0)}(\lambda), \psi^{(k)}]$$

with  $\psi^{(k)} \in \mathcal{H}_{\Lambda,+}^k(M_T)$ ,  $k = \overline{0, m+2}$ . The following theorem [3, 4] based on mappings (25) holds.

**Theorem 3.** *The set of operators (25) parametrized by  $\lambda \in \Sigma$  realizes the cohomology group isomorphism formulated in Theorem 1.2*

*Proof.* A proof of this theorem one can get passing over in (25) to the corresponding cohomology  $\mathcal{H}_{\Lambda(\mathcal{L}),+}^k(M_T)$  and homology  $H_k(M_T; \mathbb{C})$  groups of  $M_T$  for every  $k = \overline{0, m+2}$ . If one to take an element  $\psi^{(k)} := \psi^{(k)}(\mu) \in \mathcal{H}_{\Lambda(\mathcal{L}),+}^k(M_T)$ ,  $k = \overline{0, m+2}$ , solving the equation  $d_{\mathcal{L}}\psi^{(k)}(\mu) = 0$  with  $\mu \in \Sigma_k$  being some set of the related "spectral" parameters marking elements of the subspace  $\mathcal{H}_{\Lambda(\mathcal{L}),-}^k(M_T)$ , then one finds easily from (25) and identity (23) that  $dZ^{(k)}[\varphi^{(0)}(\lambda), \psi^{(k)}(\mu)] = 0$  for all  $(\lambda, \mu) \in \Sigma \times \Sigma_k$ ,  $k = \overline{0, m+2}$ . This, in particular, means due to the Poincare lemma

[27, 28, 29] that there exist differential  $(k-1)$ -forms  $\Omega^{(k-1)}[\varphi^{(0)}(\lambda), \psi^{(k)}(\mu)] \in \Lambda^{k-1}(M; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , such that

$$(26) \quad Z^{(k)}[\varphi^{(0)}(\lambda), \psi^{(k)}(\mu)] = d\Omega^{(k-1)}[\varphi^{(0)}(\lambda), \psi^{(k)}(\mu)]$$

for all pairs  $(\varphi^{(0)}(\lambda), \psi^{(k)}(\mu)) \in \mathcal{H}_0^* \times \mathcal{H}_{\Lambda(\mathcal{L}),+}^k(M_T)$  parametrized by  $(\lambda, \mu) \in \Sigma \times \Sigma_k$ ,  $k = \overline{0, m+2}$ . As a result of passing on the right hand-side of (25) to the homology groups  $H_k(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , one gets due to the standard Stokes theorem [27, 29, 28] that the mappings

$$(27) \quad B_\lambda^{(k)} : \mathcal{H}_{\Lambda(\mathcal{L}),+}^k(M_T) \longrightarrow H_k(M_T; \mathbb{C})$$

are isomorphisms for every  $k = \overline{0, m+2}$  and  $\lambda \in \Sigma$ . Making further use of the Poincare duality [8, 28, 29] between the homology groups  $H_k(M_T; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , and the cohomology groups  $H^k(M; \mathbb{C})$ ,  $k = \overline{0, m+2}$ , respectively, one obtains finally the statement claimed in Theorem 1.4.  $\triangleright$

□

## 2. THE SPECTRAL STRUCTURE OF DELSARTE-DARBOUX TYPE TRANSMUTATION

### OPERATORS IN MULTIDIMENSION

**2.1** Take now into account that our differential operators  $L_j : \mathcal{H} \rightarrow \mathcal{H}$ ,  $j = \overline{1, 2}$ , are of the special form (2). Assume also that differential expressions (4) are normal closed operators defined on dense subspace  $D(L) \subset L_2(M; \mathbb{C}^N)$ .

Then due to Theorem 1.4 one can find such a pair  $(\varphi^{(0)}(\lambda), \psi^{(k)}(\mu)) \in \mathcal{H}_0^* \times \mathcal{H}_{\Lambda(\mathcal{L}),+}^k(M_T)$  parametrized by elements  $(\lambda, \mu) \in \Sigma \times \Sigma_k$ , for which the equality

$$(28) \quad B_\lambda^{(m)}(\psi^{(0)}(\mu)dx) = S_{(t;x)}^{(m)} \int_{\partial S_{(t;x)}^{(m)}} \Omega^{(m-1)}[\varphi^{(0)}(\lambda), \psi^{(0)}(\mu)dx]$$

holds, where  $S_{(t;x)}^{(m)} \in H_m(M_T; \mathbb{C})$  is some arbitrary but fixed element of parametrized by an arbitrarily chosen point  $(t; x) \in M_T \cap \partial S_{(t;x)}^{(m)}$ . Consider the next integral expressions

$$(29) \quad \begin{aligned} \Omega_{(t;x)}^{(m-1)}(\lambda, \mu) &: = \int_{\sigma_{(t;x)}^{(m-1)}} \Omega^{(m-1)}[\varphi^{(0)}(\lambda), \psi^{(0)}(\mu)dx], \\ \Omega_{(t_0;x_0)}^{(m-1)}(\lambda, \mu) &: = \int_{\sigma_{(t_0;x_0)}^{(m-1)}} \Omega^{(m-1)}[\phi^{(0)}(\lambda), \psi^{(0)}(\mu)dx], \end{aligned}$$

with a point  $(t_0; x_0) \in M_T \cap \partial S_{(t_0;x_0)}^{(m)}$  being taken fixed the boundaries  $\sigma_{(t;x)}^{(m-1)} := \partial S_{t;x}^{(m)}$ ,  $\sigma_{(t_0;x_0)}^{(m-1)} := \partial S_{t_0;x_0}^{(m)}$  assumed to be homological to each other as  $(t; x_0) \rightarrow (t; x) \in M_T$ ,  $(\lambda, \mu) \in \Sigma \times \Sigma_k$ , and interpret them as the kernels [12, 13, 32] of the corresponding invertible integral operators of Hilbert-Schmidt type  $\Omega_{(t;x)}, \Omega_{(t_0;x_0)} : L_2^{(\rho)}(\Sigma; \mathbb{C}) \rightarrow L_2^{(\rho)}(\Sigma; \mathbb{C})$ , where  $\rho$  is some finite Borel measure on the parameters set  $\Sigma$ . Define now the invertible operators expressions

$$(30) \quad \Omega_\pm : \psi^{(0)}(\mu) \longrightarrow \tilde{\psi}^{(0)}(\mu)$$

for  $\psi^{(0)}(\mu)dx \in \mathcal{H}_{\Lambda(\mathcal{L}),+}^m(M_T)$  and some  $\tilde{\psi}^{(0)}(\mu)dx \in \mathcal{H}_{\Lambda(\mathcal{L}),+}^m(M_T)$ ,  $\mu \in \Sigma$ , where, by definition, for any  $\eta \in \Sigma$

$$(31) \quad \begin{aligned} \tilde{\psi}^{(0)}(\eta) &: = \psi^{(0)}(\eta) \cdot \Omega_{(t;x)}^{-1} \cdot \Omega_{(t_0;x_0)} \\ &= \int_{\Sigma} d\rho(\mu) \int_{\Sigma} d\rho(\xi) \psi^{(0)}(\mu) \Omega_{(t;x)}^{-1}(\mu, \xi) \Omega_{(t_0;x_0)}(\xi, \eta), \end{aligned}$$

being motivated by the expression (28). Namely, consider the following diagram

$$(32) \quad \begin{array}{ccc} \mathcal{H}_{\Lambda(\mathcal{L}),+}^m(M_T) & \xrightarrow{\Omega_{\pm}} & \mathcal{H}_{\Lambda(\tilde{\mathcal{L}}),+}^m(M_T), \\ B_{\lambda}^{(m)} \downarrow & \swarrow \tilde{B}_{\lambda}^{(m)} & \\ H_m(M_T; \mathbb{C}) & & \end{array}$$

which is assumed to be commutative for some another co-chain complex

$$(33) \quad \mathcal{H} \longrightarrow \Lambda^0(M_T; \mathcal{H}) \xrightarrow{d_{\tilde{\mathcal{L}}}} \Lambda^1(M_T; \mathcal{H}) \xrightarrow{d_{\tilde{\mathcal{L}}}} \dots \xrightarrow{d_{\tilde{\mathcal{L}}}} \Lambda^{m+2}(M_T; \mathcal{H}) \xrightarrow{d_{\tilde{\mathcal{L}}}} 0.$$

Here, by definition, the generalized anti-differentiation is

$$(34) \quad d_{\tilde{\mathcal{L}}} := \sum_{j=1}^2 dt_j \wedge \tilde{L}_j(t; x|\partial)$$

with

$$(35) \quad \begin{aligned} \tilde{L}_j &= \partial/\partial t_j - \tilde{L}_j(t; x|\partial), \\ \tilde{L}_j(t; x|\partial) &: = \sum_{|\alpha|=0}^{n_j(\tilde{L})} \tilde{a}_{\alpha}^{(j)}(t; x) \frac{\partial^{|\alpha|}}{\partial x^{\alpha}}, \end{aligned}$$

where coefficients  $\tilde{a}_{\alpha}^{(j)} \in C^1(T; C^{\infty}(M; \text{End } \mathbb{C}^N))$ ,  $|\alpha| = 0, n_j(\tilde{L})$ ,  $n_j(\tilde{L}) := n_j(L) \in \mathbb{Z}_+$ ,  $j = \overline{1, 2}$ . The corresponding isomorphisms  $\tilde{B}_{\lambda}^{(m)} : \mathcal{H}_{\Lambda(\mathcal{L}),+}^m(M_T) \longrightarrow H_m(M_T; \mathbb{C})$ ,  $\lambda \in \Sigma$ , act, by definition, as follows:

$$(36) \quad \tilde{B}_{\lambda}^{(m)}(\tilde{\psi}^{(0)}(\mu)dx) = S_{(t;x)}^{(m)} \int_{\partial S_{(t;x)}^{(m)}} \tilde{\Omega}^{(m-1)}[\tilde{\varphi}^{(0)}(\lambda), \tilde{\psi}^{(0)}(\mu)dx],$$

where  $\tilde{\varphi}^{(0)}(\lambda) \in \tilde{\mathcal{H}}_0^* \subset \mathcal{H}_{\Lambda(\mathcal{L}^*),-}^0(M_T)$ ,  $\lambda \in (\sigma(\tilde{L}) \cap \bar{\sigma}(\tilde{L}^*)) \times \Sigma_{\sigma}$ ,

$$(37) \quad \tilde{\mathcal{H}}_0^* := \{\tilde{\varphi}^{(0)}(\lambda) \in \mathcal{H}_{\Lambda(\mathcal{L}^*),-}^n(M_T) : d_{\tilde{\mathcal{L}}}^* \tilde{\varphi}^{(0)}(\lambda) = 0, \tilde{\varphi}^{(0)}(\lambda)|_{\tilde{\Gamma}} = 0, \lambda \in \Sigma\}$$

for some hypersurface  $\tilde{\Gamma} \subset M_T$ . Respectively, one defines the following closed subspace

$$(38) \quad \tilde{\mathcal{H}}_0 := \{\tilde{\psi}^{(0)}(\mu) \in \mathcal{H}_{\Lambda(\mathcal{L}^*),-}^0(M_T) : d_{\tilde{\mathcal{L}}}^* \tilde{\psi}^{(0)}(\mu) = 0, \tilde{\psi}^{(0)}(\mu)|_{\tilde{\Gamma}} = 0, \mu \in \Sigma\}$$

for the hyperspace  $\tilde{\Gamma} \subset M_T$ , introduced above.

Suppose now that the elements (31) belong to the closed subspace (38), that is

$$(39) \quad d_{\tilde{\mathcal{L}}} \tilde{\psi}^{(0)}(\mu) = 0$$

Define similarly to (38) a closed subspace  $\tilde{\mathcal{H}}_0^* \subset \mathcal{H}_{\Lambda(\mathcal{L}^*),-}^m(M_T)$  as follows:

$$(40) \quad \mathcal{H}_0 := \{\psi^{(0)}(\lambda) \in \mathcal{H}_{\Lambda(\mathcal{L}^*),-}^0(M_T) : d_{\mathcal{L}} \psi^{(0)}(\lambda) = 0, \psi^{(0)}(\lambda)|_{\Gamma} = 0, \lambda \in \Sigma\}$$

for all  $\mu \in \Sigma$ . Then due to the commutativity of the diagram (32) there exist the corresponding two invertible mappings

$$(41) \quad \Omega_{\pm} : \mathcal{H}_0 \rightarrow \tilde{\mathcal{H}}_0,$$

depending on ways of their extending over the whole Hilbert space  $\mathcal{H}_{\Lambda,-}^m(M_T)$ . Extend now operators (41) upon the whole Hilbert space  $\mathcal{H}_{\Lambda,-}^m(M_T)$  by means of the standard method [21, 23] of variation of constants, taking into account that for kernels  $\Omega_{(t;x)}(\lambda, \mu), \Omega_{(t_0;x_0)}(\lambda, \mu) \in L_2^{(p)}(\Sigma; \mathbb{C}) \otimes L_2^{(p)}(\Sigma; \mathbb{C})$ ,  $\lambda, \mu \in \Sigma$ , one can write down the following relationships:

$$(42) \quad \begin{aligned} & \Omega_{(t;x)}(\lambda, \mu) - \Omega_{(t_0;x_0)}(\lambda, \mu) \\ &= \int_{\partial S_{(t;x)}^{(m)}} \Omega^{(m-1)}[\varphi^{(0)}(x), \psi^{(0)}(\mu) dx] - \int_{\partial S_{(t_0;x_0)}^{(m)}} \Omega^{(m-1)}[\varphi^{(0)}(\lambda), \psi^{(0)}(\mu) dx] \\ &= \int_{S_{\pm}^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)})} d\Omega^{(m-1)}[\varphi^{(0)}(\lambda), \psi^{(0)}(\mu) dx] \\ &= \int_{S_{\pm}^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)})} Z^{(m)}[\varphi^{(0)}(\lambda), \psi^{(0)}(\mu) dx], \end{aligned}$$

where, by definition,  $m$ -dimensional open surfaces  $S_{\pm}^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)}) \subset M_T$  are spanned smoothly without self-intersection between two homological cycles  $\sigma_{(t;x)}^{(m-1)} = \partial S_{(t;x)}^{(m)}$  and  $\sigma_{(t_0;x_0)}^{(m-1)} = \partial S_{(t_0;x_0)}^{(m)} \in C_{m-1}(M_T; \mathbb{C})$  in such a way that the boundary  $\partial(S_{+}^{(m)}(\sigma_{(t_0;x_0)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)}) \cup S_{-}^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)})) = \emptyset$ . Making use of the relationship (42), one can thereby find easily the following integral operator expressions in  $\mathcal{H}_{-}$ :

$$(43) \quad \begin{aligned} \Omega_{\pm} &= \mathbf{1} - \int_{\Sigma} d\rho(\eta) \tilde{\psi}^{(0)}(\xi) \Omega_{(t_0;x_0)}^{-1}(\xi, \eta) \\ &\quad \times \int_{S_{\pm}^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)})} Z^{(m)}[\varphi^{(0)}(\eta), (\cdot) dx] \end{aligned}$$

defined for fixed pairs  $(\varphi^{(0)}(\xi), \psi^{(0)}(\eta)) \in \mathcal{H}_0^* \times \mathcal{H}_0$  and  $(\tilde{\varphi}^{(0)}(\xi), \tilde{\psi}^{(0)}(\mu)) \in \tilde{\mathcal{H}}_0^* \times \tilde{\mathcal{H}}_0$ ,  $\lambda, \mu \in \Sigma$ , being bounded invertible operators of Volterra type [18, 19, 14, 32] on the whole Hilbert space  $\mathcal{H}$ . Moreover, for the differential operators  $\tilde{L}_j : \mathcal{H} \rightarrow \mathcal{H}$ ,  $j = \overline{1, 2}$ , one can get easily the following expressions

$$(44) \quad \tilde{L}_j = \Omega_{\pm} L_j \Omega_{\pm}^{-1}$$

for  $j = \overline{1, 2}$ , where the left hand-side of (44) does not depend on signs " $\pm$ " of the right-hand sides. Thereby, the Volterrian integral operators (43) are the Delsarte-Darboux transmutation operators, mapping a given set  $\mathcal{L}$  of differential operators into a new set  $\tilde{\mathcal{L}}$  of differential operators transformed via the Delsarte expressions (44).



**2.2** Suppose now that all of differential operators  $L_j(t; x|\partial)$ ,  $j = \overline{1, 2}$ , considered above don't depend on the variable  $t \in T$ . Then, evidently, one can take

$$\begin{aligned}
 \mathcal{H}_0 & : = \{\psi_\mu^{(0)}(\xi) \in L_{2,-}(M; \mathbb{C}^N) : L_j \psi_\mu^{(0)}(\xi) = \mu_j \psi_\mu^{(0)}(\xi), \\
 j & = \overline{1, 2}, \psi_\mu^{(0)}(\xi)|_{\tilde{\Gamma}} = 0, \mu = (\mu_1, \mu_2) \in \sigma(\tilde{L}) \cap \overline{\sigma(L^*)}, \xi \in \Sigma_\sigma\} \\
 \tilde{\mathcal{H}}_0 & : = \{\tilde{\psi}_\mu^{(0)}(\xi) \in L_{2,-}(M; \mathbb{C}^N) : \tilde{L}_j \tilde{\psi}_\mu^{(0)}(\xi) = \mu_j \tilde{\psi}_\mu^{(0)}(\xi), \\
 j & = \overline{1, 2}, \tilde{\psi}_\mu^{(0)}(\xi)|_{\tilde{\Gamma}} = 0, \mu = (\mu_1, \mu_2) \in \sigma(\tilde{L}) \cap \overline{\sigma(L^*)}, \xi \in \Sigma_\sigma\} \\
 (45) \quad \mathcal{H}_0^* & : = \{\varphi_\lambda^{(0)}(\eta) \in L_{2,-}(M; \mathbb{C}^N) : L_j^* \varphi_\lambda^{(0)}(\eta) = \bar{\lambda}_j \varphi_\lambda^{(0)}(\eta), j = \overline{1, 2}, \\
 \varphi_\lambda^{(0)}(\eta)|_{\tilde{\Gamma}} & = 0, \lambda = (\lambda_1, \lambda_2) \in \sigma(\tilde{L}) \cap \overline{\sigma(L^*)}, \eta \in \Sigma_\sigma\} \\
 \tilde{\mathcal{H}}_0^* & : = \{\tilde{\varphi}_\lambda^{(0)}(\eta) \in L_{2,-}(M; \mathbb{C}^N) : \tilde{L}_j^* \tilde{\varphi}_\lambda^{(0)}(\eta) = \bar{\lambda}_j \tilde{\varphi}_\lambda^{(0)}(\eta), \\
 j & = \overline{1, 2}, \tilde{\varphi}_\lambda^{(0)}(\eta)|_{\tilde{\Gamma}} = 0, \lambda = (\lambda_1, \lambda_2) \in \sigma(\tilde{L}) \cap \overline{\sigma(L^*)}, \eta \in \Sigma_\sigma\}
 \end{aligned}$$

and construct the corresponding Delsarte-Darboux transmutation operators

$$\begin{aligned}
 (46) \quad \Omega_\pm & = 1 - \int_{\sigma(\tilde{L}) \cap \overline{\sigma(L^*)}} d\rho_\sigma(\lambda) \int_{\Sigma_\sigma \times \Sigma_\sigma} d\rho_{\Sigma_\sigma}(\xi) d\rho_{\Sigma_\sigma}(\eta) \\
 & \times \int_{S_\pm^{(m)} \sigma_{(t_0; x_0)}^{(m-1)}, \sigma_{(t_0; x_0)}^{(m-1)}} dx \tilde{\psi}_\lambda^{(0)}(\xi) \Omega_{x_0}^{-1}(\lambda; \xi; \eta) \tilde{\varphi}_\lambda^{(0), \top}(\eta)(\cdot)
 \end{aligned}$$

acting already in the Hilbert space  $L_{2,+}(M; \mathbb{C}^N)$ , where for any  $(\lambda; \xi, \eta) \in (\sigma(\tilde{L}) \cap \overline{\sigma(L^*)} \times \Sigma_\sigma^2)$  kernels

$$(47) \quad \Omega_{(x_0)}(\lambda; \xi, \eta) := \int_{\sigma_{x_0}^{(m-1)}} \Omega^{(m-1)}[\varphi_\lambda^{(0)}(\xi), \psi_\lambda^{(0)}(\eta) dx]$$

for  $(\xi, \eta) \in \Sigma_\sigma^2$  and every  $\lambda \in \sigma(\tilde{L}) \cap \overline{\sigma(L^*)}$  belong to  $L_2^{(\rho)}(\Sigma_\sigma; \mathbb{C}) \otimes L_2^{(\rho)}(\Sigma_\sigma; \mathbb{C})$ . Moreover, as  $\partial \Omega_\pm / \partial t_j = 0$ ,  $j = \overline{1, 2}$ , one gets easily the set of differential expressions

$$(48) \quad \tilde{L}_j(x|\partial) := \Omega_\pm L_j(x|\partial) \Omega_\pm^{-1}$$

$j = \overline{1, 2}$ , also commuting, evidently, to each other.

The Volterrian operators (46) possess some additional properties. Namely, define the following Fredholm type integral operator in  $H$  :

$$(49) \quad \Omega := \Omega_+^{-1} \Omega_-,$$

which can be written in the form

$$(50) \quad \Omega = \mathbf{1} + \Phi(\Omega),$$

where the operator  $\Phi(\Omega) \in \mathcal{B}_\infty(H)$  is compact. Moreover, due to the relationships (48) one gets easily that the following commutator conditions

$$(51) \quad [\Omega, L_j] = 0$$

hold for  $j = \overline{1, 2}$ .

Denote now by  $\hat{\Phi}(\Omega) \in H_- \otimes H_-$  and  $\hat{K}_+(\Omega), \hat{K}_-(\Omega) \in H_- \otimes H_-$  the kernels corresponding [12, 13] to operators  $\Phi(\Omega) \in \mathcal{B}_\infty(H)$  and  $\Omega_\pm - \mathbf{1} \in \mathcal{B}_\infty(H)$ . Then due to the fact that supports  $\text{supp } K_+(\Omega) \cap \text{supp } K_-(\Omega) = \sigma_x^{(m-1)} \cup \sigma_{x_0}^{(m-1)}$ , one

gets from (49) and (50) the well known Gelfand-Levitan-Marchenko linear integral equation

$$(52) \quad \hat{K}_+(\Omega) + \hat{\Phi}(\Omega) + \hat{K}_+(\Omega)_+ * \hat{\Phi}(\Omega) = \hat{K}_-(\Omega),$$

allowing to find the factorizing the Fredholmian operator (49) kernel  $\hat{K}_+(\Omega)(x; y) \in H_- \otimes H_-$  for all  $y \in \text{supp} K_+(\Omega)$ . The conditions (51) can be rewritten suitably as follows:

$$(53) \quad (L_{j,ext} \otimes \mathbf{1})\hat{\Phi}(\Omega) = (1 \otimes L_{j,ext}^*)\hat{\Phi}(\Omega)$$

for  $j = \overline{1, 2}$ , where  $L_{j,ext} \in \mathcal{L}(H_-)$ ,  $j = \overline{1, 2}$ , and their adjoint  $L_{j,ext}^* \in \mathcal{L}(H_-)$ ,  $j = \overline{1, 2}$ , are the corresponding extensions [12, 24, 13] of the differential operators  $L_j$  and  $L_j^* \in \mathcal{L}(H)$ ,  $j = \overline{1, 2}$ .

Concerning the relationships (48) one can write down [12, 24] kernel conditions similar to (53):

$$(54) \quad (\tilde{L}_{j,ext} \otimes \mathbf{1})\hat{K}_\pm(\Omega) = (\mathbf{1} \otimes L_{j,ext}^*)\hat{K}_\pm(\Omega),$$

where as above,  $\tilde{L}_{j,ext} \in \mathcal{L}(H_-)$ ,  $j = \overline{1, 2}$  are the corresponding rigging extensions of the differential operators  $\tilde{L}_j \in \mathcal{L}(H)$ ,  $j = \overline{1, 2}$ .

**2.3** Proceed now to analyzing the question about the general differential and spectral structure of transformed operator expression (44). It is evidently that found above conditions (52) and (53) on the kernels  $\hat{K}_\pm(\Omega) \in \mathcal{H}_- \otimes \mathcal{H}_-$  of Delsarte-Darboux transmutation operators are necessary for the operator expressions (44) to exist and be differential. Put the question whether these conditions are also sufficient?

For studying this question let us consider Volterrian operators (43) and (46) with kernels satisfying the conditions (52) and (53), assuming that suitable oriented surfaces  $S_\pm^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)}) \in C_m(M_T; \mathbb{C})$  are given as follows:

$$(55) \quad \begin{aligned} S_+^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)}) &= \{(t'; x') \in M_T : \\ &\quad t' = P(t; x|x'), t \in T\}, \\ S_-^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)}) &= \{(t'; x') \in M_T : \\ &\quad t' = P(t; x|x') \in T \setminus [t_0, t]\}, \end{aligned}$$

where a mapping  $P \in C^\infty(M_T \times M; T)$  is smooth and such that the boundaries  $\partial S_\pm^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)}) = \pm(\sigma_{(t;x)}^{(m-1)} - \sigma_{(t_0;x_0)}^{(m-1)})$  with cycles  $\sigma_{(t;x)}^{(m-1)}$  and  $\sigma_{(t_0;x_0)}^{(m-1)} \in \mathcal{K}(M_T)$  being homological to each other for any choice of points  $(t_0; x_0)$  and  $(t; x) \in M_T$ . Then one can see by means of some simple but cumbersome calculations, based on considerations from [34] and [9], that the resulting expressions on the right hand-sides of

$$(56) \quad \tilde{L} = L + [K_\pm(\Omega), L] \cdot \Omega_\pm^{-1}$$

are exactly equal to each other differential ones if such there was the expression for an operator  $L \in \mathcal{L}(\mathcal{H})$ .

Concerning the inverse operators  $\Omega_\pm^{-1} \in \mathcal{B}(\mathcal{H})$  present in (56) one can notice here that due to the functional symmetry between closed subspaces  $\mathcal{H}_0$  and  $\tilde{\mathcal{H}}_0 \subset$

$\tilde{\mathcal{H}}_-$ , the defining relationships (41) and (31) are reversible, that is there exist the inverse operator mappings  $\Omega_{\pm}^{-1} : \tilde{\mathcal{H}}_0 \rightarrow \mathcal{H}_0$ , such that

$$(57) \quad \Omega_{\pm}^{-1} : \tilde{\psi}^{(0)}(\lambda) \longrightarrow \psi^{(0)}(\lambda) := \tilde{\psi}^{(0)}(\lambda) \cdot \tilde{\Omega}_{(t;x)}^{-1} \tilde{\Omega}_{(t;x)}$$

for some suitable kernels  $\tilde{\Omega}_{(t;x)}(\lambda, \mu)$  and  $\tilde{\Omega}_{(t_0;x_0)}(\lambda, \mu) \in L_2^{(\rho)}(\Sigma; \mathbb{C}) \otimes L_2^{(\rho)}(\Sigma; \mathbb{C})$ , related naturally with the transformed differential expression  $\tilde{L} \in \mathcal{L}(\mathcal{H})$ . Thereby, due to the expressions (57) one can write down similar to (46) the following inverse integral operators:

$$(58) \quad \begin{aligned} \Omega_{\pm}^{-1} &= \mathbf{1} - \int_{\Sigma} d\rho(\xi) \int_{\Sigma} d\rho(\eta) \psi^{(0)}(\xi) \tilde{\Omega}_{t_0;x_0}^{-1}(\xi, \eta) \\ &\quad \times \int_{S_{\pm}^{(m)}(\sigma_{(t;x)}^{(m-1)}, \sigma_{(t_0;x_0)}^{(m-1)})} \tilde{Z}^{(m)}[\tilde{\varphi}^{(0)}(\eta), (\cdot) dx] \end{aligned}$$

defined for fixed pairs  $(\tilde{\varphi}^{(0)}(\xi), \tilde{\psi}^{(0)}(\eta)) \in \tilde{\mathcal{H}}_0^* \times \tilde{\mathcal{H}}_0$  and  $(\varphi^{(0)}(\xi), \psi^{(0)}(\eta)) \in \mathcal{H}_0^* \times \mathcal{H}_0$ ,  $\xi, \eta \in \Sigma$ , and being bounded invertible operators of Volterra type in the whole Hilbert space  $\mathcal{H}$ . In particular, the compatibility conditions  $\Omega_{\pm} \Omega_{\pm}^{-1} = \mathbf{1} = \Omega_{\pm}^{-1} \Omega_{\pm}$  must be fulfilled identically in  $\mathcal{H}$ , involving some restrictions identifying measures  $\rho$  and  $\Sigma$  and possible asymptotic conditions of coefficient functions of the differential expression  $L \in \mathcal{L}$ . Such kinds of restrictions were already mentioned before in [37, 38, 39], where in particular the relationships with the local and nonlocal Riemann problems were discussed.

**2.4** Within the framework of the general construction presented above one can give a natural interpretation of so called Backlund transformations for coefficient functions of a given differential operator expression  $L \in \mathcal{L}(\mathcal{H})$ . Namely, following the symbolic considerations in [41], we reinterpret the approach devised there for constructing the Backlund transformations making use of the techniques based on the theory of Delsarte transmutation operators. Let us define two different Delsarte-Darboux transformed differential operator expressions

$$(59) \quad L_1 = \Omega_{1,\pm} L \Omega_{1,\pm}^{-1}, \quad L_2 = \Omega_{2,\pm} L \Omega_{2,\pm}^{-1},$$

where  $\Omega_{1,+}, \Omega_{2,-} \in \mathcal{B}(\mathcal{H})$  are some Delsarte transmutation Volterrian operators in  $\mathcal{H}$  with Borel spectral measures  $\rho_1$  and  $\rho_2$  on  $\Sigma$ , such that the following conditions

$$(60) \quad \Omega_{1,+}^{-1} \Omega_{1,-} = \Omega = \Omega_{2,+}^{-1} \Omega_{2,-}$$

hold. Making use now of the conditions (59) and relationships (60) one finds easily that the operator  $B := \Omega_{2,-} \Omega_{1,+}^{-1} \in \mathcal{B}(\mathcal{H})$  satisfies the following operator equations:

$$(61) \quad L_2 B = B L_1, \quad \Omega_{2,\pm} B = B \Omega_{1,\pm},$$

which motivate the next definition.

**Definition 4.** *An invertible symbolic mapping  $B : \mathcal{L}(\mathcal{H}) \longrightarrow \mathcal{L}(\mathcal{H})$  will be called a Darboux-Backlund transformation of an operator  $L_1 \in \mathcal{L}(\mathcal{H})$  into the operator  $L_2 \in \mathcal{L}(\mathcal{H})$  if there holds the condition*

$$(62) \quad [QB, L_1] = 0$$

for some linear differential expression  $Q \in \mathcal{L}(\mathcal{H})$ .

The condition (62) can be realized as follows. Take any differential expression  $q \in \mathcal{L}(\mathcal{H})$  satisfying the symbolic equation

$$(63) \quad [qB, L] = 0.$$

Then, making use of the transformations like (59), from (60) one finds that

$$(64) \quad [QB, L_1] = 0,$$

where owing to (61)

$$(65) \quad QB := \Omega_{1,+} q B \Omega_{1,+}^{-1} = \Omega_{1,+} q \Omega_{2,+}^{-1} B.$$

Therefore, the expression  $Q = \Omega_{1,+} q \Omega_{2,+}^{-1}$  appears to be differential one too owing to the conditions (61).

The consideration above related with the symbolic mapping  $B : \mathcal{L}(\mathcal{H}) \rightarrow \mathcal{L}(\mathcal{H})$  gives rise to an effective tool of constructing self-Backlund transformations for coefficients of differential operator expressions  $L_1, L_2 \in \mathcal{L}(\mathcal{H})$  having many applications [15, 35, 26, 32, 23] in spectral and soliton theories.

**2.5** Return now back to studying the structure Delsarte-Darboux transformations for a polynomial differential operators pencil

$$(66) \quad L(\lambda; x|\partial) := \sum_{j=0}^{n(L)} L_j(x|\partial) \lambda^j,$$

where  $n(L) \in \mathbb{Z}_+$  and  $\lambda \in \mathbb{C}$  is a complex-valued parameter. It is asked to find the corresponding to (66) Delsarte-Darboux transformations  $\Omega_{\lambda,\pm} \in \mathcal{B}(\mathcal{H})$ ,  $\lambda \in \mathbb{C}$ , such that for some polynomial differential operators pencil  $\tilde{L}(\lambda; x|\partial) \in \mathcal{L}(\mathcal{H})$  the following Delsarte-Lions [2] transmutation condition

$$(67) \quad \tilde{L} \Omega_{\lambda,\pm} = \Omega_{\lambda,\pm} L$$

holds for almost all  $\lambda \in \mathbb{C}$ . For such transformations  $\Omega_{\lambda,\pm} \in \mathcal{B}(\mathcal{H})$  to be found, let us consider a parameter  $\tau \in \mathbb{R}$  dependent differential operator  $L_\tau(x|\partial) \in \mathcal{L}(\mathcal{H}_\tau)$ , where

$$(68) \quad L_\tau(x|\partial) := \sum_{j=0}^{n(L)} L_j(x|\partial) \partial^j / \partial \tau^j,$$

acting in the functional space  $\mathcal{H}_\tau = C^{q(L)}(\mathbb{R}_\tau; \mathcal{H})$  for some  $q(L) \in \mathbb{Z}_+$ . Then one can easily construct the corresponding Delsarte-Darboux transformations  $\Omega_{\tau,\pm} \in \mathcal{B}(\mathcal{H}_\tau)$  of Volterra type for some differential operator expression

$$(69) \quad \tilde{L}_\tau(x|\partial) := \sum_{j=0}^{n(L)} \tilde{L}_j(x|\partial) \partial^j / \partial \tau^j,$$

if the following Delsarte-Lions [2] transmutation conditions

$$(70) \quad \tilde{L}_\tau \Omega_{\tau,\pm} = \Omega_{\tau,\pm} L_\tau$$

hold in  $\mathcal{H}_\tau$ . Thus, making use of the results obtained above, one can write down that

$$(71) \quad \begin{aligned} \Omega_{\tau,\pm} &= \mathbf{1} - \int_{\Sigma} d\rho_{\Sigma}(\xi) \int_{\Sigma} d\rho_{\Sigma}(\eta) \tilde{\psi}_{\tau}^{(0)}(\lambda; \xi) \Omega_{(\tau_0; x_0)}^{-1}(\lambda; \xi, \eta) \\ &\quad \times \int_{S_{\pm}^{(m)}(\sigma_{(\tau; x)}^{(m-1)}, \sigma_{(\tau_0; x_0)}^{(m-1)})} Z^{(m)}[\varphi_{\tau}^{(0)}(\lambda; \eta), (\cdot) dx] \end{aligned}$$

defined by means of the following closed subspaces  $\mathcal{H}_{\tau,0} \subset \mathcal{H}_{\tau,-}$  and  $\mathcal{H}_{\tau,0}^* \subset \mathcal{H}_{\tau,-}^*$  :

$$\begin{aligned} \mathcal{H}_{\tau,0} &: = \{\psi_{\tau}^{(0)}(\lambda; \xi) \in \mathcal{H}_{\tau,-} : L_{\tau} \psi_{\tau}^{(0)}(\lambda; \xi) = 0, \\ \psi_{\tau}^{(0)}(\lambda; \xi)|_{\tau=0} &= \psi^{(0)}(\lambda; \xi) \in \mathcal{H}, L \psi^{(0)}(\lambda; \xi) = 0, \\ \psi^{(0)}(\lambda; \xi)|_{\Gamma} &= 0, \lambda \in \mathbb{C}, \xi \in \Sigma\}, \end{aligned}$$

$$(72) \quad \begin{aligned} \mathcal{H}_{\tau,0}^* &: = \{\varphi_{\tau}^{(0)}(\lambda; \eta) \in \mathcal{H}_{\tau,-}^* : L_{\tau} \varphi_{\tau}^{(0)}(\lambda; \eta) = 0, \\ \varphi_{\tau}^{(0)}(\lambda; \eta)|_{\tau=0} &= \varphi^{(0)}(\lambda; \eta) \in \mathcal{H}^*, L \varphi^{(0)}(\lambda; \eta) = 0, \\ \varphi^{(0)}(\lambda; \eta)|_{\Gamma} &= 0, \lambda \in \mathbb{C}, \eta \in \Sigma\}. \end{aligned}$$

Recalling now that our operators  $L_j \in \mathcal{L}(\mathcal{H})$ ,  $j = \overline{0, r(L)}$ , do not depend on the parameter  $\tau \in \mathbb{R}$ , one can derive easily from (71)

$$(73) \quad \begin{aligned} \Omega_{\pm} &= \mathbf{1} - \int_{\Sigma} d\rho_{\Sigma}(\xi) \int_{\Sigma} d\rho_{\Sigma}(\eta) \tilde{\psi}^{(0)}(\lambda; \xi) \Omega_{(x_0)}^{-1}(\lambda; \xi, \eta) \\ &\quad \times \int_{S_{\pm}^{(m)}(\sigma_{(x)}^{(m-1)}, \sigma_{(x_0)}^{(m-1)})} Z_0^{(m)}[\varphi^{(0)}(\lambda; \eta), (\cdot) dx], \end{aligned}$$

where we put  $\sigma_x^{(m-1)} := \sigma_{(\tau_0; x)}^{(m-1)}$ ,  $\sigma_{x_0}^{(m-1)} := \sigma_{(\tau_0; x_0)}^{(m-1)} \in C_{m-1}(\mathbb{R}^m; \mathbb{C})$  and

$$(74) \quad Z_0^{(m)}[\varphi^{(0)}(\lambda; \eta), \psi^{(0)} dx] := Z^{(m)}[\varphi_{\tau}^{(0)}(\lambda; \eta), \psi_{\tau}^{(0)} dx]|_{d\tau=0}.$$

The corresponding to (73) closed subspaces  $\mathcal{H}_0 \in \mathcal{H}_-$  and  $\mathcal{H}_0^* \in \mathcal{H}_-^*$  are given as

follows:

$$(75) \quad \mathcal{H}_0 := \{\psi^{(0)}(\lambda; \xi) \in \mathcal{H}_- : L \psi^{(0)}(\lambda; \xi) = 0, \psi^{(0)}(\lambda; \xi)|_{\Gamma} = 0, \lambda \in \mathbb{C}, \xi \in \Sigma\},$$

$$\mathcal{H}_{\tau,0}^* := \{\varphi^{(0)}(\lambda; \eta) \in \mathcal{H}_-^* : L \varphi^{(0)}(\lambda; \eta) = 0, \varphi^{(0)}(\lambda; \eta)|_{\Gamma} = 0, \lambda \in \mathbb{C}, \eta \in \Sigma\}.$$

Thereby, making use of the expressions (73) one can construct the Delsarte-Darboux transformed linear differential pencil  $\tilde{L} \in \mathcal{L}(\mathcal{H})$ , whose coefficients are related with those of the pencil  $L \in \mathcal{L}(\mathcal{H})$  via some Backlund type relationships useful for applications (see [23, 20, 42, 43, 38]) in the soliton theory.

### 3. DELSARTE-DARBOUX TYPE TRANSMUTATION OPERATORS FOR SPECIAL MULTI-DIMENSIONAL EXPRESSIONS AND THEIR APPLICATIONS

#### 3.1 A perturbed self-adjoint Laplace operator in $\mathbb{R}^n$ . Consider the Laplace

operator  $-\Delta_m$  in  $H := L(\mathbb{R}^m; \mathbb{C})$  perturbed by the multiplication operator on a function  $q \in W_2^2(\mathbb{R}^m; \mathbb{C})$ , that is the operator

$$(76) \quad L(x|\partial) := -\Delta_m + q(x),$$

where  $x \in \mathbb{R}^m$ . The operator (76) is self-adjoint in  $H$ . Applying the results from the Section 1 to the differential expression (76) in the Hilbert space  $H$ , one can write down the following invertible Delsarte-Darboux transmutation operators:

$$(77) \quad \Omega_{\pm} = \mathbf{1} - \int_{\sigma(L)} d\rho_{\sigma}(\xi) \int_{\sigma(L)} d\rho_{\sigma}(\xi) \int_{\Sigma_{\sigma}} d\rho_{\Sigma_{\sigma}}(\xi) \int_{\Sigma_{\sigma}} d\rho_{\Sigma_{\sigma}}(\eta) \\ \times \tilde{\psi}^{(0)}(\lambda; \xi) \Omega_{(x_0)}^{-1}(\lambda; \xi, \eta) \int_{S_{\pm}^{(m)}(\sigma_x^{(m-1)}, \sigma_{(x_0)}^{(m-1)})}^{(0)} dy \bar{\varphi}^{(0)\top}(\lambda; \eta), (\cdot),$$

where  $\sigma_x^{(m-1)} \in \mathcal{K}(\mathbb{R}^m)$  is some closed maybe non-compact simplicial hypersurface in  $\mathbb{R}^m$  parametrized by a running point  $x \in \sigma_x^{(m-1)}$ , and  $\sigma_{x_0}^{(m-1)} \in \mathcal{K}(\mathbb{R}^m)$

is a suitable homological to  $\sigma_x^{(m-1)}$  simplicial hypersurface in  $\mathbb{R}^m$  parametrized by a point  $x_0 \in \sigma_{x_0}^{(m-1)}$ . There exist exactly two  $m$ -dimensional subspaces spanning them, say  $S_{\pm}^{(m)}(\sigma_x^{(m-1)}, \sigma_{x_0}^{(m-1)}) \in \mathcal{K}(\mathbb{R}^m)$ , such that  $S_{+}^{(m)}(\sigma_x^{(m-1)}, \sigma_{x_0}^{(m-1)}) \cup S_{-}^{(m)}(\sigma_x^{(m-1)}, \sigma_{x_0}^{(m-1)}) = \mathbb{R}^m$ . Taking into account these subspaces, one can rewrite down compactly the Delsarte-Darboux transmutation operators (77) for (76):

$$(78) \quad \Omega_{\pm} = \mathbf{1} + \int_{S_{\pm}^{(m)}(\sigma_x^{(m-1)}, \sigma_{x_0}^{(m-1)})} dy \hat{K}_{\pm}(\Omega)(x; y)(\cdot),$$

where, as before,  $x \in \sigma_x^{(m-1)}$  and kernels  $\hat{K}_{\pm}(\Omega) \in H_{-} \otimes H_{-}$  satisfy the equations (54), or equivalently,

$$(79) \quad -\Delta_m(x; \partial) \hat{K}_{\pm}(\Omega)(x; y) + \Delta_m(y; \partial) \hat{K}_{\pm}(\Omega)(x; y) \\ = (q(y) - \tilde{q}(x)) \hat{K}_{\pm}(\Omega)(x; y)$$

for all  $x, y \in \text{supp } \hat{K}_{\pm}(\Omega)$ . Take for simplicity, a non-compact closed simplicial

hypersurface  $\sigma_x^{(m-1)} = \sigma_{x, \gamma}^{(m-1)} := \{y \in \mathbb{R}^m : \langle x - y, \pm \gamma \rangle = 0\}$  and the degenerate simplicial cycle  $\sigma_{x_0}^{(m-1)} := x_0 = \infty \in \mathbb{R}^m$ , where  $\gamma \in \mathbb{S}^{m-1}$  is an arbitrary versor,  $||\gamma|| = 0$ . Then, evidently,

$$(80) \quad S_{\pm}^{(m)}(\sigma_{x, \gamma}^{(m-1)}, \sigma_{\infty}^{(m-1)}) := S_{\pm \gamma, x}^{(m)} = \{y \in \mathbb{R}^m : \langle x - y, \pm \gamma \rangle \geq 0\}$$

and our transmutation operators (78) take the form

$$(81) \quad \Omega_{\pm \gamma} = \mathbf{1} + \int_{S_{\pm \gamma, x}^{(m)}} dy \hat{K}_{\pm \gamma}(\Omega)(x; y)(\cdot),$$

where  $\text{supp } \hat{K}_{\pm \gamma}(\Omega) = S_{\pm \gamma, x}^{(m)}, S_{+ \gamma, x}^{(m)} \cap S_{- \gamma, x}^{(m)} = \sigma_{x, \gamma}^{(m-1)} \cup \sigma_{\infty}^{(m-1)}$  and  $S_{+ \gamma, x}^{(m)} \cap S_{- \gamma, x}^{(m)} = \mathbb{R}^m$  for any direction  $\gamma \in \mathbb{S}^{m-1}$ .

The invertible transmutation Volterrian operators like (81) were constructed before by L.D. Faddeev [9] for the self-adjoint perturbed Laplace operator (76)

in  $\mathbb{R}^3$ . He called them [9] transformation operators with a Volterrian direction  $\gamma \in \mathbb{S}^{m-1}$ . It is easy to see that Faddeev's expressions (81) are very special cases of the general expressions (78) obtained above.

Define now making use of (78) the following Fredholmian operator in the Hilbert space  $H$  :

$$(82) \quad \mathbf{\Omega} := (\mathbf{1} + K_+(\mathbf{\Omega}))^{-1}(\mathbf{1} + K_-(\mathbf{\Omega})) = \mathbf{1} + \Phi(\mathbf{\Omega})$$

with the compact part  $\Phi(\mathbf{\Omega}) \in \mathcal{B}_\infty(H)$ . Then the commutation equality

$$(83) \quad [L, \Phi(\mathbf{\Omega})] = \mathbf{0}$$

together with the Gelfand-Levitan-Marchenko equation

$$(84) \quad K_+(\mathbf{\Omega}) + \hat{\Phi}(\mathbf{\Omega}) + \hat{K}_+(\mathbf{\Omega}) + \hat{\Phi}(\mathbf{\Omega}) = \hat{K}_-(\mathbf{\Omega})$$

for the corresponding kernels  $\hat{K}_\pm(\mathbf{\Omega})$  and  $\hat{\Phi}(\mathbf{\Omega}) \in H_- \otimes H_-$  hold.

In [9] there was thoroughly analyzed the spectral structure of kernels  $\hat{K}_\pm(\mathbf{\Omega}) \in H_- \otimes H_-$  in (81) making use of the analytical properties of the corresponding Green functions of the operator (76). As one can see from (77), these properties depend strongly both on the structure of the spectral measures  $\rho_\sigma$  on  $\sigma(L)$  and  $\rho_{\Sigma_\sigma}$  on  $\Sigma_\sigma$  and on analytical behavior of the kernel  $\Omega_\infty(\lambda; \xi, \eta) \in L_2^{(\rho)}(\Sigma_\sigma; \mathbb{C}) \otimes L_2^{(\rho)}(\Sigma_\sigma; \mathbb{C})$ ,  $\xi, \eta \in \Sigma_\sigma$ , for all  $\lambda \in \sigma(L)$ . In [9] there was stated for any direction  $\gamma \in \mathbb{S}^{m-1}$  the dependence of kernels  $\hat{K}_\pm(\mathbf{\Omega}) \in H_- \otimes H_-$  on the regularized determinant of the resolvent  $R_\mu(L) \in \mathcal{B}(H)$ ,  $\mu \in \mathbb{C}/\sigma(L)$  is a regular point, for the operator (76). This dependence can be also clarified if to make use of the approach from Section 2.

**2.2 A two-dimensional Dirac type operator.** Let us define in  $H := L_2(\mathbb{R}^2; \mathbb{C}^2)$  a two-dimensional Dirac type operator

$$(85) \quad \tilde{L}(x; \partial) := \begin{pmatrix} \partial/\partial x_1 & \tilde{u}_1(x) \\ \tilde{u}_2(x) & \partial/\partial x_2 \end{pmatrix},$$

where  $x := (x_1, x_2) \in \mathbb{R}^2$ , and coefficients  $\tilde{u}_j \in W_2^1(\mathbb{R}^2; \mathbb{C})$ ,  $j = \overline{1, 2}$ . the transformation properties of the operator (85) were studied [16] thoroughly by L.P. Nizhnik. In particular, he constructed some special class of the Delsarte-Darboux transmutation operators in the form

$$(86) \quad \mathbf{\Omega}_\pm = \mathbf{1} + \int_{S_\pm^{(2)}(\sigma_x^{(1)}, \sigma_\infty^{(1)})} dy \hat{K}_\pm(\mathbf{\Omega})(x; y)(\cdot),$$

where for two orthonormal versors  $\gamma_1$  and  $\gamma_2 \in \mathbb{S}^1$ ,  $||\gamma_1|| = 1 = ||\gamma_2||$ ,

$$(87) \quad \begin{aligned} S_+^{(2)}(\sigma_x^{(1)}, \sigma_\infty^{(1)}) & : = \{y \in \mathbb{R}^2 : \langle x - y, \gamma_1 \rangle \geq 0\} \\ & \cap \{y \in \mathbb{R}^2 : \langle x - y, \gamma_2 \rangle \geq 0\}, \\ S_-^{(2)}(\sigma_x^{(1)}, \sigma_\infty^{(1)}) & : = \{y \in \mathbb{R}^2 : \langle x - y, \gamma_1 \rangle \leq 0\} \\ & \cup \{y \in \mathbb{R}^2 : \langle x - y, \gamma_2 \rangle \leq 0\}. \end{aligned}$$

In the case when  $\langle x, \gamma_j \rangle = x_j \in \mathbb{R}$ ,  $j = \overline{1, 2}$ , the corresponding kernel

$$(88) \quad \hat{K}_+(\mathbf{\Omega}) = \begin{pmatrix} K_{+,11}^{(1)} \delta_{\langle y-x, \gamma_1 \rangle} + K_{+,11}^{(0)}(x; y) & K_{+,12}^{(1)} \delta_{\langle y-x, \gamma_2 \rangle} + K_{+,12}^{(0)}(x; y) \\ K_{+,21}^{(1)} \delta_{\langle y-x, \gamma_1 \rangle} + K_{+,21}^{(0)}(x; y) & K_{+,22}^{(1)} \delta_{\langle y-x, \gamma_2 \rangle} + K_{+,22}^{(0)}(x; y) \end{pmatrix}$$

is Dirac delta-function singular, being, in part, localized on half-lines  $< y - x, \gamma_2 > = 0$  and  $< y - x, \gamma_1 > = 0$ , with regular all coefficients  $K_{+,ij}^{(l)} \in C^1(\mathbb{R}^2 \times \mathbb{R}^2; \mathbb{C})$  for all  $i, j = \overline{1, 2}$  and  $l = \overline{0, 1}$ . Such a property of the transmutation kernels for the perturbed Laplace operator (76) was also observed in [9], where it was motivated by the necessary condition for the transformed operator  $\tilde{L}(x; \partial) \in \mathcal{L}(H)$  to

be differential. As one can check, the same reason of the existence of singularities holds in (88).

Let us now consider the general expression like (78) for the corresponding hyper-surfaces  $S_{\pm}^{(2)}(\sigma_x^{(1)}, \sigma_{\infty}^{(1)})$  spanning between a closed non-compact smooth cycle  $\sigma_x^{(1)} \in \mathcal{K}(\mathbb{R}^2)$  and the infinite point  $\sigma_{\infty}^{(1)} := \infty \in \mathcal{K}(\mathbb{R}^2)$ . A running point  $x \in \sigma_x^{(1)}$  is taken arbitrary but, as usual, fixed. the kernels  $\hat{K}_{\pm}(\Omega) \in H_- \times H_-$  in (86) satisfy

the standard conditions (53) and (54), that is

$$(89) \quad \begin{aligned} (\tilde{L}_{1,ext} \otimes \mathbf{1}) \hat{K}_{\pm}(\Omega) &= (\mathbf{1} \otimes L_{1,ext}^*) \hat{K}_{\pm}(\Omega), \\ [L_1, \Phi(\Omega)] &= 0 \end{aligned}$$

for some matrix differential Dirac type operator  $L_1 \in \mathcal{L}(H)$  of the form (76). Together with this Dirac operator the following matrix second order differential operator

$$(90) \quad \tilde{L}_2 := \mathbf{1} \frac{\partial}{\partial t} + \begin{pmatrix} \frac{\partial^2}{\partial x_1^2} \pm \frac{\partial^2}{\partial x_2^2} - \tilde{v}_2 & -2 \frac{\partial \tilde{u}_1}{\partial x_2} \\ -2 \frac{\partial \tilde{u}_2}{\partial x_1} & \frac{\partial^2}{\partial x_1^2} \pm \frac{\partial^2}{\partial x_2^2} - \tilde{v}_1 \end{pmatrix}$$

in the parametric space  $\mathcal{H} := C^1(\mathbb{R}; H)$  was studied in [16, 17] for which there was

developed scattering theory and given its application for constructing soliton-like exact solutions to the so called Davey-Stewartson nonlinear dynamical system in partial derivatives. The latter was based on the fact that two operators  $\tilde{L}_1$  and  $\tilde{L}_2 \in \mathcal{L}(H)$  are commuting to each other.

Namely, consider the Volterrian operators  $\Omega_{\pm} \in \mathcal{B}(\mathcal{H})$  realizing the following Delsarte-Darboux transmutations:

$$(91) \quad \tilde{L}_1 \Omega_{\pm} = \Omega_{\pm} L_1, \quad \tilde{L}_2 \Omega_{\pm} = \Omega_{\pm} L_2.$$

Here we put

$$(92) \quad \begin{aligned} L_1(x; \partial) &: = \begin{pmatrix} \partial/\partial x_1 & 0 \\ 0 & \partial/\partial x_2 \end{pmatrix}, \\ L_2 &: = \mathbf{1} \frac{\partial}{\partial t} + \begin{pmatrix} \frac{\partial^2}{\partial x_1^2} \pm \frac{\partial^2}{\partial x_2^2} - \alpha_2(x_2) & 0 \\ 0 & \frac{\partial^2}{\partial x_1^2} \pm \frac{\partial^2}{\partial x_2^2} - \alpha_1(x_1) \end{pmatrix}, \end{aligned}$$

where  $\alpha_j \in W_2^1(\mathbb{R}; \mathbb{C})$ ,  $j = \overline{1, 2}$ , are some given functions. It is evident that operators (92) are commuting to each other. then, if the operators  $\Omega_{\pm} \in \mathcal{B}(H)$  exist and satisfy (91), the following commutation condition

$$(93) \quad [\tilde{L}_1, \tilde{L}_2] = 0$$



holds, that there was exactly claimed above and effectively exploited before in [16, 17].

Recall now that for the operators  $\mathbf{\Omega}_\pm \in \mathcal{B}(H)$  to exist they must satisfy additionally the kernel conditions (89) and

$$(94) \quad \begin{aligned} (\tilde{L}_{2,ext} \otimes \mathbf{1}) \hat{K}_\pm(\mathbf{\Omega}) &= (\mathbf{1} \otimes L_{2,ext}^*) \hat{K}_\pm(\mathbf{\Omega}), \\ [L_2, \Phi(\mathbf{\Omega})] &= 0, \end{aligned}$$

where, as before, the operator  $\Phi(\mathbf{\Omega}) \in \mathcal{B}_\infty(H)$  is defined by (82) as

$$(95) \quad \mathbf{\Omega} := \mathbf{1} + \Phi(\mathbf{\Omega}).$$

Owing to the evident commutation condition (93) the set of equations (89) and (94) is compatible giving rise to the expression like (86), where the kernel  $\hat{K}_+(\mathbf{\Omega}) \in H_- \otimes H_-$  satisfies the set of differential equations generalizing those from [16, 17]:

$$(96) \quad \begin{aligned} \frac{\partial K_{+,11}}{\partial x_1} + \frac{\partial K_{+,11}}{\partial y_1} + \tilde{u}_1 K_{+,21} &= 0, \quad \frac{\partial K_{+,12}}{\partial x_1} + \frac{\partial K_{+,12}}{\partial y_1} + \tilde{u}_1 K_{+,22} = 0, \\ \frac{\partial K_{+,21}}{\partial x_2} + \frac{\partial K_{+,21}}{\partial x_1} + \tilde{u}_2 K_{+,11} &= 0, \quad \frac{\partial K_{+,22}}{\partial x_2} + \frac{\partial K_{+,22}}{\partial y_2} + \tilde{u}_2 K_{+,12} = 0, \\ \pm \frac{\partial \tilde{u}_1}{\partial x_2} K_{+,21} &= \frac{\partial K_{+,11}}{\partial t} + \left[ \left( \frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial y_1^2} \right) \pm \left( \frac{\partial^2}{\partial x_2^2} - \frac{\partial^2}{\partial y_2^2} \right) \right] K_{+,11} \\ &\quad + (\alpha_2(x_2) - \tilde{v}_2(x)) K_{+,11} \\ \pm \frac{\partial \tilde{u}_1}{\partial x_2} K_{+,21} &= \frac{\partial K_{+,22}}{\partial t} + \left[ \left( \frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial y_1^2} \right) \pm \left( \frac{\partial^2}{\partial x_2^2} - \frac{\partial^2}{\partial y_2^2} \right) \right] K_{+,22} \\ &\quad + (\alpha_1(x_1) - \tilde{v}_1(x)) K_{+,22}, \\ \mp 2 \frac{\partial \tilde{u}_1}{\partial x_2} K_{+,22} &= \frac{\partial K_{+,12}}{\partial t} + \left[ \left( \frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial y_1^2} \right) \pm \left( \frac{\partial^2}{\partial x_2^2} - \frac{\partial^2}{\partial y_2^2} \right) \right] K_{+,12} \\ &\quad + (\alpha_1(x_1) - \tilde{v}_2(x)) K_{+,22}, \\ 2 \frac{\partial \tilde{u}_2}{\partial x_1} K_{+,22} &= \frac{\partial K_{+,21}}{\partial t} + \left[ \left( \frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial y_1^2} \right) \pm \left( \frac{\partial^2}{\partial x_2^2} - \frac{\partial^2}{\partial y_2^2} \right) \right] K_{+,21} \\ &\quad + (\alpha_2(x_2) - \tilde{v}_1(x)) K_{+,11}. \end{aligned}$$

Moreover, the following conditions

$$(97) \quad \begin{aligned} \tilde{u}_1(x) &= -K_{+,12}^{(0)}|_{y=x}, \quad \tilde{u}_2(x) = -K_{+,21}^{(0)}|_{y=x}, \\ \tilde{v}_2(x)|_{x_1=-\infty} &= \alpha_2(x_2), \quad \tilde{v}_1(x)|_{x_2=-\infty} = \alpha_1(x_1) \end{aligned}$$

hold for all  $x \in \mathbb{R}^2$  and  $y \in \text{supp} \hat{K}_+(\mathbf{\Omega})$ , where we take into account the singular series expansion

$$(98) \quad \hat{K}_+(\mathbf{\Omega}) = \sum_{s=0}^{p(K_+)} K_+^{(s)} \delta_{\sigma_x^{(1)}}^{(s-1)}$$

for some finite integer  $p(K_+) \in \mathbb{Z}_+$  with respect to the Dirac function  $\delta_{\sigma_x^{(1)}} : W_2^q(\mathbb{R}^2; \mathbb{C}) \rightarrow \mathbb{R}$ ,  $q \in \mathbb{Z}_+$ , and its derivatives, having the support (see [34], Chapter

3) coinciding with the closed cycle  $\sigma_x^{(1)} \in \mathcal{K}(\mathbb{R}^2)$ .

**Remark 5.** Concerning the special case (88) discussed before in [16, 17], one gets easily that  $p(K_+) = 1$  and  $\sigma_x^{(1)} = \partial(\cap_{j=\overline{1,2}}\{y \in \mathbb{R}^2 : \langle y-x, \gamma_j \rangle = 0\}) \subset \text{supp} \hat{K}_+(\Omega)$ . It was shown also before that equations like (96) and (97) possess solutions if the Gelfand-Levitan-Marchenko equation (52) does.

Making use also of the exact forms of operators  $L_1$  and  $L_2 \in \mathcal{L}(\mathcal{H})$ , one obtains easily from (89) and (94) the corresponding set of differential equations for components of the kernel  $\hat{\Phi}(\Omega) \in H_- \otimes H_-$  :

$$(99) \quad \begin{aligned} \frac{\partial \Phi_{11}}{\partial x_1} + \frac{\partial \Phi_{11}}{\partial y_1} &= 0, & \frac{\partial \Phi_{12}}{\partial x_1} + \frac{\partial \Phi_{12}}{\partial y_1} &= 0, \\ \frac{\partial \Phi_{21}}{\partial x_2} + \frac{\partial \Phi_{21}}{\partial y_2} &= 0, & \frac{\partial \Phi_{22}}{\partial x_2} + \frac{\partial \Phi_{22}}{\partial y_2} &= 0, \\ \frac{\partial \Phi_{11}}{\partial t} \pm (\frac{\partial^2}{\partial x_2^2} - \frac{\partial^2}{\partial y_2^2})\Phi_{11} + (\alpha_2(y_2) - \alpha_2(x_2))\Phi_{11} &= 0, \\ \frac{\partial \Phi_{12}}{\partial t} \pm (\frac{\partial^2}{\partial x_2^2} - \frac{\partial^2}{\partial y_2^2})\Phi_{12} + (\alpha_1(y_1) - \alpha_2(x_2))\Phi_{12} &= 0, \\ \frac{\partial \Phi_{21}}{\partial t} + (\frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial y_1^2})\Phi_{21} + (\alpha_2(y_2) - \alpha_1(x_1))\Phi_{21} &= 0, \\ \frac{\partial \Phi_{22}}{\partial t} + (\frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial y_1^2})\Phi_{22} + (\alpha_1(y_1) - \alpha_1(x_1))\Phi_{22} &= 0 \end{aligned}$$

for all  $(x, y) \in \mathbb{R}^2 \times \mathbb{R}^2$ . The obtained above equations (99) generalize those before found in [16, 17] and used for exact integrating the well known Devey-Stewartson differential equation [37, 35, 10] and finding so called soliton like solutions. Concerning our generalized case the kernel (98) is a solution to the following Gelfand-Levitan-Marchenko type equations:

$$(100) \quad \begin{aligned} K_+^{(0)}(x; y) + \Phi^{(0)}(x; y) + \int_{S_+^{(2)}(\sigma_x^{(1)}, \sigma_\infty^{(1)})} K_+^{(0)}(x; \xi) \Phi^{(0)}(\xi; y) d\xi \\ + \int_{\sigma_x^{(1)}} K_+^{(1)}(x; \xi) \Phi^{(0)}(\xi; y) d\sigma_x^{(1)} = 0, \\ K_+^{(1)}(x; y) + \Phi^{(1)}(x; y) + \int_{S_+^{(2)}(\sigma_x^{(1)}, \sigma_\infty^{(1)})} K_+^{(0)}(x; \xi) \Phi^{(1)}(\xi; y) d\xi \\ + \int_{\sigma_x^{(1)}} K_+^{(1)}(x; \xi) \Phi^{(1)}(\xi; y) d\sigma_x^{(1)} = 0, \end{aligned}$$

where  $y \in S_+^{(2)}(\sigma_x^{(1)}, \sigma_\infty^{(1)})$  for all  $x \in \mathbb{R}^2$  and, by definition,

$$(101) \quad \hat{\Phi}(\Omega) := \Phi^{(0)} + \Phi^{(1)} \delta_{\sigma_x^{(1)}}$$

is the corresponding to (98) kernel expansion. Since the kernel (101) is singular, the differential equations (99) must be treated naturally in the distributional sense [34].

Taking into account the exact forms of "dressed" differential operators  $L_j \in \mathcal{L}(\mathcal{H})$ ,  $j = \overline{1, 2}$ , given by (85) and (90) one gets easily that the commutativity condition (93) gives rise to that of  $\tilde{L}_j \in \mathcal{L}(\mathcal{H})$ ,  $j = \overline{1, 2}$ , being equivalent to the mentioned before Devey-Stewartson dynamical system

$$\begin{aligned}
(102) \quad d\tilde{u}_1/dt &= -(\tilde{u}_{1,xx} + \tilde{u}_{1,yy}) + 2(\tilde{v}_1 - \tilde{v}_2), \\
d\tilde{u}_2/dt &= \tilde{u}_{2,xx} + \tilde{u}_{2,yy} + 2(\tilde{v}_2 - \tilde{v}_1), \\
\tilde{v}_{1,x} &= (\tilde{u}_1\tilde{u}_2)_y, \quad \tilde{v}_{2,x} = (\tilde{u}_1\tilde{u}_2)_x
\end{aligned}$$

on a functional infinite-dymensioanl manifold  $M_u \subset \mathcal{S}(\mathbb{R}^2; \mathbb{C})$ . The exact soliton like solutions to (102) are given by expressions (97), where the kernel  $K_+^{(1)}(\Omega)$  solves the second linear integral equation of (100). On the other hand-side, there exists the exact expression (31) which solves the set of "dressed" equations

$$(103) \quad \tilde{L}_1 \tilde{\psi}^{(0)}(\eta) = 0, \quad \tilde{L}_2 \tilde{\psi}^{(0)}(\eta) = 0.$$

Since the kernels  $\Omega(\lambda, \mu) \in L_2^{(\rho)}(\Sigma; \mathbb{C}) \otimes L_2^{(\rho)}(\Sigma; \mathbb{C})$ , for  $\lambda, \mu \in \Sigma$ ,  $(t; x) \in M_T \cap S_+^{(2)}(\sigma_x^{(1)}, \sigma_\infty^{(1)})$  are given by means of exact expressions (29), one can find via simple calculations the corresponding analytical expression for the functions  $(\tilde{u}_1, \tilde{u}_2) \in M_u$ , solving the dynamical system (102). This procedure is often called the Darboux type transformation and was recently extensively used as a particular case of the construction above in [23] for finding soliton-like solutions to the Devey-Stewartson (102) and related with it two-dimensional modified Korteweg-de Vries flows on  $M_u$ . Moreover, as it can be observed from the technique used for constructing the Delsarte-Darboux transmutation operators  $\Omega_\pm \in \mathcal{B}(\mathcal{H})$ , the set of solutions to (102) obtained by means of Darboux type transformations coincides

completely with the corresponding set of solutions obtained by means of solving the related set of Gelfand-Levitan-Marchenko integral equations (99) and (100).

**2.3 An affine de Rham-Hodge-Skrypnik differential complex and related generalized self-dual Yang-Mills flows.** Consider the following set of affine differential expressions in  $\mathcal{H} := C^1(\mathbb{R}^{m+1}; H)$ ,  $H := L_2(\mathbb{R}^m; \mathbb{C}^N)$  :

$$(104) \quad L_i(\lambda) := \mathbf{1} \frac{\partial}{\partial p_i} - \lambda \frac{\partial}{\partial x_i} + A_i(x; p|t),$$

where  $x \in \mathbb{R}^m$ ,  $(t, p) \in \mathbb{R}^{m+1}$ , matrices  $A_i \in C^1(\mathbb{R}^{m+1}; S(\mathbb{R}^m; \text{End} \mathbb{C}^N))$ ,  $i = \overline{1, m}$ , and a parameter  $\lambda \in \mathbb{C}$ . One can easily now construct an exact affine de Rham-Hodge-Skrypnik differential complex on  $M_T := \mathbb{R}^{m+1} \times \mathbb{R}^m$  as

$$(105) \quad \mathcal{H} \rightarrow \Lambda(M_T; \mathcal{H}) \xrightarrow{d_{\mathcal{L}(\lambda)}} \Lambda^1(M_T; \mathcal{H}) \xrightarrow{d_{\mathcal{L}(\lambda)}} \Lambda^2(M_T; \mathcal{H}) \xrightarrow{d_{\mathcal{L}(\lambda)}} 0,$$

where, by definition, the differentiation

$$(106) \quad d_{\mathcal{L}(\lambda)} := dt \wedge B(\lambda) + \sum_{i=1}^m dp_i \wedge L_i(\lambda)$$

and the affine matrix

$$(107) \quad B(\lambda) := \partial/\partial t - \sum_{s=0}^{n(B)+q} B_s(x; p|t) \lambda^{n(B)-s}$$

with matrices  $B_s \in C^1(\mathbb{R}^{m+1}; S(\mathbb{R}^m; \text{End} \mathbb{C}^N))$ ,  $s = \overline{0, n(B)+q}$ ,  $n(B), q \in \mathbb{Z}_+$ . The affine complex (105) will be exact for all  $\lambda \in \mathbb{C}$  iff the following generalized self-dual Yang-Mills equations [43]

$$\partial A_i / \partial p_j - \partial A_j / \partial p_i - [A_i, A_j] = 0, \quad \partial A_i / \partial x_j - \partial A_j / \partial x_i = 0,$$

$$(108) \quad \begin{aligned} \partial B_0 / \partial x_i &= 0, \quad \partial B_{n(B)+q} / \partial p_i = 0, \quad \partial B_s / \partial x_i = \partial B_{s-1} / \partial p_i + [A_i, B_{s-1}] = 0, \\ \partial A_i / \partial t + \partial B_{n(B)} / \partial p_i - \partial B_{n(B)+1} / \partial x_i + [A_i, B_{n(B)}] &= 0 \end{aligned}$$

hold for all  $i, j = \overline{1, m}$  and  $s = \overline{0, n(B) \vee n(B) + q, n(B) + 2}$ . Assume now that the conditions (108) are satisfied on  $M_T$ . Then, making the change  $\mathbb{C} \ni \lambda \rightarrow \partial / \partial \tau : \mathcal{H} \rightarrow \mathcal{H}$ ,  $\tau \in \mathbb{R}$ , one finds the following set of pure differential expressions

$$(109) \quad \begin{aligned} L_{i(\tau)} &: = \mathbf{1} \frac{\partial}{\partial p_i} - \frac{\partial^2}{\partial \tau \partial x_i} + A_i(x; p|t), \\ B_{(\tau)} &: = \partial / \partial t - \sum_{s=0}^{n(B)+q} B_s(x; p|t) \left( \frac{\partial}{\partial \tau} \right)^{n(B)-s}, \end{aligned}$$

where matrices  $A_i$ ,  $i = \overline{1, m}$ , and  $B_s$ ,  $s = \overline{0, n(B) + q}$ , don't depend on the variable  $\tau \in \mathbb{R}$ . By means operator expressions (109) one can now naturally construct a new differential complex related with that of (105):

$$(110) \quad \mathcal{H}_{(\tau)} \rightarrow \Lambda(M_{T,\tau}; \mathcal{H}_{(\tau)}) \xrightarrow{d_{\mathcal{L}}} \Lambda^1(M_{T,\tau}; \mathcal{H}_{(\tau)}) \rightarrow \dots \rightarrow \Lambda^{2m+2}(M_{T,\tau}; \mathcal{H}_{(\tau)}) \xrightarrow{d_{\mathcal{L}}} 0,$$

where, by definition,  $\mathcal{H}_{(\tau)} := C^1(\mathbb{R}^{m+1}; H_{(\tau)})$ ,  $H_{(\tau)} := L_2(\mathbb{R}^m \times \mathbb{R}_\tau; \mathbb{C}^N)$  and

$$(111) \quad d_{\mathcal{L}} := dt \wedge B_{(\tau)} + \sum_{i=1}^m dp_i \wedge L_{i(\tau)}.$$

Owing to the condition (108) the following lemma holds.

**Lemma 6.** *The differential complex (110) is exact.*

Therefore, one can build the standard de Rham-Hodge-Skrypnik type Hilbert space decomposition

$$(112) \quad \mathcal{H}_\Lambda(M_{T,\tau}) := \bigoplus_{k=0}^{k=2m+2} \mathcal{H}_\Lambda^k(M_{T,\tau})$$

as well the corresponding Hilbert-Schmidt rigging

$$(113) \quad \mathcal{H}_{\Lambda,+}(M_{T,\tau}) \subset \mathcal{H}_\Lambda(M_{T,\tau}) \subset \mathcal{H}_{\Lambda,-}(M_{T,\tau}).$$

Making use now of the results obtained in subsection 1.5, one can define the Delsarte closed subspaces  $\mathcal{H}_{0(\tau)}$  and  $\tilde{\mathcal{H}}_{0(\tau)} \subset \mathcal{H}_{(\tau)-}$ , related with the exact complex (110):

$$(114) \quad \begin{aligned} \mathcal{H}_{0(\tau)} &: = \{ \psi_{(\tau)}^{(0)}(\xi) \in \mathcal{H}_{\Lambda,-}^0(M_{T,\tau}) : L_{j(\tau)} \psi_{(\tau)}^{(0)}(\xi) = 0, \\ B_{(\tau)} \psi_{(\tau)}^{(0)}(\xi) &= 0, \quad \psi_{(\tau)}^{(0)}(\xi)|_\Gamma = 0, \quad \psi_{(\tau)}^{(0)}(\xi)|_{t=0} = e^{\lambda\tau} \psi_\lambda^{(0)}(\eta) \in \mathcal{H}_{\Lambda,-}^0(M_{\mathbb{R}^m,\tau}), \\ L_j(\lambda) \psi_\lambda^{(0)}(\eta) &= 0, \quad \xi = (\lambda; \eta) \in \Sigma := \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)} \}, \end{aligned}$$

$$\begin{aligned} \tilde{\mathcal{H}}_{0(\tau)} &: = \{ \tilde{\psi}_{(\tau)}^{(0)}(\xi) \in \mathcal{H}_{\Lambda,-}^0(M_{T,\tau}) : \tilde{L}_{j(\tau)}^{(0)} \tilde{\psi}_{(\tau)}^{(0)}(\xi) = 0, \\ \tilde{B}_{(\tau)} \tilde{\psi}_{(\tau)}^{(0)}(\xi) &= 0, \quad \tilde{\psi}_{(\tau)}^{(0)}(\xi)|_{\tilde{\Gamma}} = 0, \quad \tilde{\psi}_{(\tau)}^{(0)}(\xi)|_{t=0} = e^{\lambda\tau} \tilde{\psi}_\lambda^{(0)}(\eta) \in \mathcal{H}_{\Lambda,-}^0(M_{\mathbb{R}^m,\tau}), \\ \tilde{L}_j(\lambda) \tilde{\psi}_\lambda^{(0)}(\eta) &= 0, \quad \xi = (\lambda; \eta) \in \Sigma := \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)} \}, \end{aligned}$$

where  $\Gamma$  and  $\tilde{\Gamma} \subset M_{T,\tau}$  are some smooth hyper-surfaces. The similar expressions

correspond to the adjoint closed subspaces  $\mathcal{H}_{0(\tau)}^*$  and  $\tilde{\mathcal{H}}_{0(\tau)}^* \subset \mathcal{H}_{\tau,-}^*$  :

$$(115) \quad \begin{aligned} \tilde{\mathcal{H}}_{0(\tau)} & : = \{\varphi_{(\tau)}^{(0)}(\xi) \in \mathcal{H}_{\Lambda,-}^0(M_{T,\tau}) : L_{j(\tau)}^* \varphi_{(\tau)}^{(0)}(\xi) = 0, \\ B_{(\tau)} \varphi_{(\tau)}^{(0)}(\xi) & = 0, \varphi_{(\tau)}^{(0)}(\xi)|_{\Gamma} = 0, \varphi_{(\tau)}^{(0)}(\xi)|_{t=0} = e^{-\bar{\lambda}\tau} \varphi_{\lambda}^{(0)}(\eta) \in \mathcal{H}_{\Lambda,-}^0(M_{\mathbb{R}^m,\tau}), \\ L_j^*(\lambda) \varphi_{\lambda}^{(0)}(\eta) & = 0, \xi = (\lambda; \eta) \in \Sigma := \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)}\}, \end{aligned}$$

$$\begin{aligned} \tilde{\mathcal{H}}_{0(\tau)} & : = \{\tilde{\varphi}_{(\tau)}^{(0)}(\xi) \in \mathcal{H}_{\Lambda,-}^0(M_{T,\tau}) : \tilde{L}_{j(\tau)}^* \tilde{\varphi}_{(\tau)}^{(0)}(\xi) = 0, \\ \tilde{B}_{(\tau)} \tilde{\varphi}_{(\tau)}^{(0)}(\xi) & = 0, \tilde{\varphi}_{(\tau)}^{(0)}(\xi)|_{\tilde{\Gamma}} = 0, \tilde{\varphi}_{(\tau)}^{(0)}(\xi)|_{t=0} = e^{-\bar{\lambda}\tau} \tilde{\varphi}_{\lambda}^{(0)}(\eta) \in \mathcal{H}_{\Lambda,-}^0(M_{\mathbb{R}^m,\tau}), \\ \tilde{L}_j^*(\lambda) \tilde{\varphi}_{\lambda}^{(0)}(\eta) & = 0, \xi = (\lambda; \eta) \in \Sigma := \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)}\}. \end{aligned}$$

Based on the closed subspaces (115) and (114), one can suitably build the Darboux type kernel  $\tilde{\Omega}_{(t,x;\tau)}(\eta, \xi) \in L_2^{(\rho)}(\Sigma_{\mathbb{C}}^{(m)}; \mathbb{C}) \otimes L_2^{(\rho)}(\Sigma_{\mathbb{C}}^{(m)}; \mathbb{C})$ ,  $\eta, \xi \in \Sigma_{\mathbb{C}}^{(m)}$ , and further, the corresponding Delsarte transmutation mappings  $\tilde{\Omega}_{\pm} \in \mathcal{B}(H_{(\tau)})$ . Namely, assume that the following conditions

$$(116) \quad \psi_{(\tau)}^{(0)}(\xi) := \tilde{\psi}_{(\tau)}^{(0)}(\xi) \cdot \tilde{\Omega}_{(t,p;x;\tau)}^{-1} \tilde{\Omega}_{(t_0,p_0,x_0;\tau)}$$

for any  $\xi \in \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)}$  hold, where

$$(117) \quad \begin{aligned} \tilde{\Omega}_{(t,x;\tau)}^{(2m+1)}(\mu, \xi) & := \int_{\sigma(t;x;\tau)} \tilde{\Omega}_{(\tau)}^{(2m+1)}[e^{-\bar{\lambda}\tau} \tilde{\varphi}^{(0)}(\mu), e^{\lambda\tau} \tilde{\psi}^{(0)}(\eta) dx \wedge dp \wedge dt], \\ \tilde{\Omega}_{(\tau)}^{(2m+1)}[e^{-\bar{\lambda}\tau} \tilde{\varphi}^{(0)}(\mu), \sum_{i=1}^m e^{\lambda\tau} \tilde{\psi}^{(0)}(\xi_{(i)}) \wedge d\tau \wedge dx \wedge \bigwedge_{j \neq i}^m dp_j] \\ & : = d\tilde{\Omega}_{(\tau)}^{(2m)}[e^{-\bar{\lambda}\tau} \tilde{\varphi}^{(0)}(\mu), \sum_{i=1}^m e^{\lambda\tau} \tilde{\psi}^{(0)}(\xi_{(i)}) \wedge d\tau \wedge dx \wedge \bigwedge_{j \neq i}^m dp_j], \end{aligned}$$

and, similarly to (24), there holds the relationship

$$(118) \quad \begin{aligned} & < d_{\tilde{\mathcal{L}}}^* \tilde{\varphi}^{(0)}(\mu) e^{-\bar{\lambda}\tau}, * \sum_{i=1}^m e^{\lambda\tau} \tilde{\psi}^{(0)}(\xi_{(i)}) dt \wedge d\tau \wedge dx \wedge \bigwedge_{j \neq i}^m dp_j > \\ & = < (*)^{-1} \tilde{\varphi}^{(0)}(\mu) e^{-\bar{\lambda}\tau}, d_{\tilde{\mathcal{L}}}(\sum_{i=1}^m e^{\lambda\tau} \tilde{\psi}^{(0)}(\xi_{(i)}) dt \wedge d\tau \wedge dx \wedge \bigwedge_{j \neq i}^m dp_j) > \\ & \quad + d\tilde{\Omega}_{(\tau)}^{(2m+1)}[\tilde{\varphi}^{(0)}(\mu) e^{-\bar{\lambda}\tau}, \sum_{i=1}^m e^{\lambda\tau} \tilde{\psi}^{(0)}(\xi_{(i)}) dt \wedge d\tau \wedge dx \wedge \bigwedge_{j \neq i}^m dp_j], \end{aligned}$$

defining the exact  $(2m+1)$ -form  $\tilde{Z}_{(\tau)}^{(2m+1)} \in \Lambda^{2m+1}(M_{T,\tau}; \mathbb{C})$ . Compute now the Delsarte transformed differential expressions

$$(119) \quad L_{j(\tau)} := \hat{\Omega}_{(\tau)\pm}^{-1} \tilde{L}_{j(\tau)} \hat{\Omega}_{(\tau)\pm}, \quad B_{(\tau)} := \hat{\Omega}_{(\tau)\pm}^{-1} \tilde{B}_{(\tau)} \hat{\Omega}_{(\tau)\pm}$$

for any  $j = \overline{1, m}$ , where, by definition,

$$(120) \quad \begin{aligned} \tilde{L}_{j(\tau)} &: = \mathbf{1} \frac{\partial}{\partial p_j} - \frac{\partial^2}{\partial \tau \partial x_j} + \bar{A}_j, \\ B_{(\tau)} &: = \partial / \partial t - \sum_{s=0}^{n(B)+q} \bar{B}_s \left( \frac{\partial}{\partial \tau} \right)^{n(B)-s} \end{aligned}$$

with all matrices  $\bar{A}_j \in \text{End} \mathbb{C}^m$ ,  $j = \overline{1, m}$ , and  $\bar{B}_s \in \text{End} \mathbb{C}^m$ ,  $s = \overline{0, n(B)+q}$ , being constant. This means, in particular, the commuting relationships

$$(121) \quad [\tilde{L}_{j(\tau)}, \tilde{L}_{i(\tau)}] = 0, \quad [\tilde{L}_{j(\tau)}, \tilde{B}_{(\tau)}] = 0$$

hold for all  $i, j = \overline{1, m}$ . Owing to the expressions (119) the induced commuting relationships

$$(122) \quad [L_{j(\tau)}, L_{i(\tau)}] = 0, \quad [L_{j(\tau)}, B_{(\tau)}] = 0$$

evidently hold, coinciding exactly with relationships (108). Moreover, reducing our differential expressions (119) upon functional subspaces  $\mathcal{H}_{(\lambda)} := e^{\lambda \tau} \mathcal{H}$ ,  $\lambda \in \mathbb{C}$ , one gets easily the set of affine differential expressions (104) and (107). Write down now the respectively reduced Delsarte transmutation operators

$$(123) \quad \begin{aligned} \hat{\Omega}_{\pm} &= \mathbf{1} - \int_{\Sigma_{\mathbb{C}}^{(m)}} d\rho_{\Sigma_{\mathbb{C}}^{(m)}}(\nu) \int_{\Sigma_{\mathbb{C}}^{(m)}} d\rho_{\Sigma_{\mathbb{C}}^{(m)}}(\eta) \psi^{(0)}(\lambda; \nu) \tilde{\Omega}_{(t_0, p_0; x_0)}^{-1}(\lambda; \nu, \eta) \\ &\times \int_{S_{\pm}^{(2m+1)}(\sigma_{(t, p; x)}^{(2m)}, \sigma_{(tt_0, p_0; x_0)}^{(2m)})} \tilde{Z}^{(2m+1)}[\tilde{\varphi}^{(0)}(\lambda; \nu), (\cdot) \sum_{i=1}^m dt \wedge dx \bigwedge_{j \neq i}^m dp_j], \end{aligned}$$

where  $\sigma_{(t, p; x)}^{(2m)}$  and  $\sigma_{(tt_0, p_0; x_0)}^{(2m)} \in \mathcal{K}(M_T)$  are some  $2m$ -dimensional closed singular simplexes, and by definition,

$$(124) \quad \begin{aligned} &\tilde{Z}^{(2m+1)}[\tilde{\varphi}^{(0)}(\lambda; \nu), \sum_{i=1}^m \tilde{\psi}^{(0)}(\lambda; \eta_{(i)}) dt \wedge dx \bigwedge_{j \neq i}^m dp_j] \\ &: = \tilde{Z}_{(\tau)}^{(2m+1)}[e^{-\bar{\lambda} \tau} \tilde{\varphi}^{(0)}(\lambda; \nu), \sum_{i=1}^m e^{\lambda \tau} \tilde{\psi}^{(0)}(\lambda; \eta_{(i)}) d\tau \wedge dt \wedge dx \bigwedge_{j \neq i}^m dp_j] \big|_{d\tau=0}, \\ &d\tilde{\Omega}_{(t, p; x)}(\lambda; \nu, \eta) := \tilde{Z}^{(2m+1)}[\tilde{\varphi}^{(0)}(\lambda; \nu), \sum_{i=1}^m \tilde{\psi}^{(0)}(\lambda; \eta_{(i)}) dt \wedge dx \bigwedge_{j \neq i}^m dp_j], \end{aligned}$$

since the  $(2m+1)$ -form (124) is owing to (118) also exact for any  $(\lambda; \nu, \eta) \in \mathbb{C} \times (\mathcal{A}_{\mathbb{C}}^{(m)} \times \mathcal{A}_{\mathbb{C}}^{(m)})$ . Thus, the operator expression (123) if applied to the operators (120) reduced upon the functional subspace  $\mathcal{H}_{(\lambda)} \simeq \mathcal{H}$ ,  $\lambda \in \mathbb{C}$ , gives rise to the differential expressions

$$(125) \quad L_j(\lambda) := \hat{\Omega}_{\pm}^{-1} \tilde{L}_j(\lambda) \hat{\Omega}_{\pm} \quad B(\lambda) := \hat{\Omega}_{\pm}^{-1} \tilde{B}(\lambda) \hat{\Omega}_{\pm},$$

where  $L_j(\lambda) \mathcal{H}_{(\lambda)} = L_{j(\tau)} \mathcal{H}_{(\lambda)}$ ,  $B(\lambda) \mathcal{H}_{(\lambda)} = B_{(\tau)}(\lambda) \mathcal{H}_{(\lambda)}$ ,  $j = \overline{1, m}$ , coinciding with affine differential expressions (104) and (107). Concerning application of these results to finding exact soliton like solutions to self-dual Yang-Mills equations (108), it is enough to mention that the relationship (116) reduced upon the subspace  $\mathcal{H}_{(\lambda)} \simeq \mathcal{H}$ ,  $\lambda \in \mathbb{C}$ , gives rise the following mapping:

$$(126) \quad \psi^{(0)}(\lambda; \eta) := \tilde{\psi}^{(0)}(\lambda; \eta) \cdot \tilde{\Omega}_{(t, p; x)}^{-1} \tilde{\Omega}_{(t_0, p_0; x_0)},$$

where kernels  $\tilde{\Omega}_{(t,p;x;\tau)}(\lambda; \eta, \xi) \in L_2^{(\rho)}(\Sigma_{\mathbb{C}}^{(m)}; \mathbb{C}) \otimes L_2^{(\rho)}(\Sigma_{\mathbb{C}}^{(m)}; \mathbb{C})$ ,  $\eta, \xi \in \Sigma_{\mathbb{C}}^{(m)}$ , for all  $(t, p; x) \in M_T$  and  $\lambda \in \mathbb{C}$ . Since the element  $\psi^{(0)}(\lambda; \eta) \in \mathcal{H}_-$  for any  $(\lambda; \xi) \in \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)}$  satisfies the set of differential equations

$$(127) \quad L_i(\lambda)\psi^{(0)}(\lambda; \eta) = 0, \quad B(\lambda)\psi^{(0)}(\lambda; \eta) = 0,$$

for all  $i = \overline{1, m}$ , from (126) and (127) one finds easily exact expressions for the corresponding matrices  $A_j$  and  $B_s \in C^1(\mathbb{R} \times \mathbb{R}^{m+1}; S(\mathbb{R}^m; \text{End } \mathbb{C}^N))$ ,  $j = \overline{1, m}$ ,  $s = \overline{0, n(B) + q}$ , satisfying the self-dual Yang-Mills equations (108). Thereby, the following theorem is stated.

**Theorem 7.** *The integral expressions (123) in  $\mathcal{H}$  are the Delsarte transmutation operators corresponding to the affine differential expressions (104), (108) and constant operators*

$$(128) \quad \tilde{L}_i(\lambda) := \mathbf{1} \frac{\partial}{\partial p_i} - \lambda \frac{\partial}{\partial x_i} + \bar{A}, \quad \tilde{B}(\lambda) := \partial / \partial t - \sum_{s=0}^{n(B)+q} \bar{B}_s \lambda^{n(B)-s}$$

for any  $\lambda \in \mathbb{C}$ . The mapping (126) realizes the isomorphisms between the closed subspaces

$$(129) \quad \begin{aligned} \mathcal{H}_0 & : = \{ \psi^{(0)}(\lambda; \eta) \in \mathcal{H}_- : d_{\tilde{\mathcal{L}}(\lambda)} \psi^{(0)}(\lambda; \eta) = 0, \psi^{(0)}(\lambda; \eta)|_{t=0} \\ & = \psi_{\lambda}^{(0)}(\eta) \in H_-, \psi^{(0)}(\lambda; \eta)|_{\Gamma} = 0, (\lambda; \eta) \in \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)} \} \end{aligned}$$

and

$$(130) \quad \begin{aligned} \tilde{\mathcal{H}}_0 & : = \{ \tilde{\psi}^{(0)}(\lambda; \eta) \in \mathcal{H}_- : d_{\tilde{\mathcal{L}}(\lambda)}^{(0)} \tilde{\psi}(\lambda; \eta) = 0, \tilde{\psi}^{(0)}(\lambda; \eta)|_{t=0} \\ & = \tilde{\psi}_{\lambda}^{(0)}(\eta) \in H_-, \tilde{\psi}^{(0)}(\lambda; \eta)|_{\tilde{\Gamma}} = 0, (\lambda; \eta) \in \mathbb{C} \times \Sigma_{\mathbb{C}}^{(m)} \} \end{aligned}$$

for any parameter  $\lambda \in \mathbb{C}$ . Moreover, the expressions (126) generate the standard Darboux type transformations for the set of operators (128) and (104), (107) via the corresponding set of linear equations (127), thereby producing exact soliton-like solutions to the self-dual Yang-Mills equations (108).

As a simple partial consequence from Theorem 3.2 one retrieves all of results obtained before in [43], where the Delsarte-Darboux mapping (126) was chosen completely a priori without any proof and motivation in the form of some affine gauge transformation.

The results similar to the above can be with a minor change applied also to the affine differential de Rham-Hodge-Skrypnik complex (105) with the external differentiation (106), where

$$(131) \quad \begin{aligned} L_i(\lambda) & : = \mathbf{1} \frac{\partial}{\partial p_i} - \left( \sum_{k=0}^{n_i(L)} a_{ik} \lambda^{k+1} \right) \frac{\partial}{\partial x_i} + \sum_{k=0}^{n_i(L)} A_{ik} \lambda^k, \\ \tilde{B}(\lambda) & : = \partial / \partial t - \sum_{s=0}^{n(B)+q} \bar{B}_s \lambda^{n(B)-s}, \end{aligned}$$

or

$$(132) \quad \begin{aligned} L_i(\lambda) &: = \mathbf{1} \frac{\partial}{\partial p_i} - \left( \sum_{k=0}^{n_i(L)} a_{ik}^{(j)} \lambda^{k+1} \right) \frac{\partial}{\partial x_j} + \sum_{k=0}^{n_i(L)} A_{ik} \lambda^k, \\ \tilde{B}(\lambda) &: = \partial / \partial t - \sum_{s=0}^{n(B)+q} \bar{B}_s \lambda^{n(B)-s}, \end{aligned}$$

for  $i = \overline{1, m}$ ,  $\lambda \in \mathbb{C}$ . The case (131) was analyzed recently in [42] by means of the

same affine gauge transformation that there was used before in [43]. To the regret, the obtained there results are too complicated and unwieldy, thereby one needs to use more mathematically motivated, clear and less cumbersome techniques for finding Delsarte-Darboux transformations and related with them soliton-like exact solutions.

#### 4. ACKNOWLEDGEMENTS

The authors thank the Organizing Committee of the 36-th Symposium on Mathematical Physics held June 9-12, 2004 in Torun University, Poland, for invitation to deliver the results of the article for Symposium audience. One of authors (A.P.) cordially thanks Prof. I.V. Skrypnyk (IM, Kyiv and IAM, Donetsk) for fruitful discussions of some aspects of the De Rham-Hodge-Skrypnyk theory and its applications presented in the article during his Kyiv-city seminar "Nonlinear Analysis" (April 14, 2004). The authors are appreciated very much to Profs. L.P. Nizhnik (IM of NAS, Kyiv), Holod P.I. (UKMA, Kyiv), T. Winiarska (IM, Politechnical University, Krakow), A. Pelczar and J. Ombach (Jagiellonian University, Krakow), J. Janas (Institute of Mathematics of PAN, Krakow), Z. Peradzynski (Warsaw University, Warsaw) and D.L. Blackmore (NJ Institute of Technology, Newark, NJ, USA) for valuable comments on diverse problems related with results presented in the article. The last but not least thanks is addressed to our friends Profs V.V. Gafiyshuk (IAPMM, Lviv) and Ya. V. Mykytiuk (I.Ya. Franko National University, Lviv) for the permanent support and help in editing the article.

#### REFERENCES

- [1] Delsarte J. *Sur certaines transformations fonctionnelles relative aux equations lineaires aux derives partielles du second ordre*. C.R. Acad. Sci. Paris, 1938, v. 206, p. 178-182
- [2] Delsarte J. and Lions J. *Transmutations d'operateurs differentielles dans le domain complex*. Comment. Math. Helv., 1957, v. 52, p. 113-128
- [3] Skrypnyk I.V. *Periods of A-closed forms*. Proceedings of the USSR Academy of Sciences, 1965, v. 160, N4, p. 772-773 (in Russian)
- [4] Skrypnyk I.V. *A harmonique fields with peculiarities*. Ukr. Math. Journal, 1965, v.17, N4, p. 130-133 (in Russian)
- [5] Skrypnyk I.V. *The generalized De Rham theorem*. Proceed. of UkrSSR Acad. of Sci., 1965, N1, p. 18-19. (in Ukrainian)
- [6] Skrypnyk I.V. *A harmonic forms on a compact Riemannian space*. Proceed. of UkrSSR Acad. of Sci., N2, p.174-175 (in Ukrainian)
- [7] Lopatynski Y.B. *On harmonic fields on Riemannian manifolds*. Ukr. Math. Journal, 1950, v.2, p. 56-60 (in Russian)
- [8] Chern S.S. *Complex manifolds*. Chicago University Publ., USA, 1956
- [9] Faddeev L.D. *Quantum inverse scattering problem. II*. in Modern problems of mathematics, M: VINITI Publ., 1974, v.3, p. 93-180 (in Russian)



- [10] Faddeev L.D. and Takhtadjan L.A. *Hamiltonian approach to soliton theory*. Moscow, Nauka, 1986 (in Russian)
- [11] Novikov S.P. (Editor) *Theory of solitons*. Moscow, Nauka Publ., 1980 (in Russian)
- [12] Berezansky Yu. M. *Eigenfunctions expansions related with selfadjoint operators*. Kiev, Nauk.Dumka Publ., 1965 (in Russian)
- [13] Berezin F.A. and Shubin M.A. *Schrodinger equation*. Moscow, the Moscow University Publisher, 1983 (in Russian)
- [14] Bukhgeim A.L. *Volterra equations and inverse problems*. Moscow, Nauka, 1983. (in Russian)
- [15] Matveev V.B. and Salle M.I. *Darboux-Backlund transformations and applications*. NY, Springer, 1993.
- [16] Nizhnik L.P. *Inverse scattering problems for hyperbolic equations*. Kiev, Nauk. Dumka Publ., 1991 (in Russian)
- [17] Nizhnik L.P. and Pochynaiko M.D. *The integration of a spatially two-dimensional Schrodinger equation by the inverse problem method*. Func. Anal. and Appl., 1982, v. 16, N1, p. 80-82 (in Russian)
- [18] Gokhberg I.C. and Krein M.G. *Theory of Volterra operators in Hilbert spaces and its applications*. Moscow, Nauka, 1967 (in Russian)
- [19] Mykytiuk Ya. V. *Factorization of Fredholmian operators*. Mathematical Studii, Proceedings of Lviv Mathematical Society, 2003, v. 20, N2, p. 185-199 (in Ukrainian)
- [20] Samoilenko A.M., Prykarpatsky Y.A. and Samoylenko V.G. *The structure of Darboux-type binary transformations and their applications in soliton theory*. Ukr. Mat. Zhurnal, 2003, v. 55, N12, p.1704-1723 (in Ukrainian)
- [21] Y.A.Prykarpatsky A.K., Samoilenko A.M. and Prykarpatsky Y.A. *The multi-dimensional Delsarte transmutation operators, their differential-geometric structure and applications. Part.1*. Opuscula Mathematica, 2003, v. 23, p.71-80 / arXiv:math-ph/0403054 v1 29 Mar 2004/
- [22] Golenia J., Prykarpatsky Y.A., Samoilenko A.M. and Prykarpatsky A.K. *The general differential-geometric structure of multidimensional Delsarte transmutation operators in parametric functional spaces and their applications in soliton theory. Part 2*. Opuscula Mathematica, 2004, N 24 /arXiv: math-ph/0403056 v 1 29 Mar 2004/
- [23] Samoilenko A.M. and Prykarpatsky Y.A. *Algebraic-analytic aspects of completely integrable dynamical systems and their perturbations*. Kyiv, NAS, Inst. Mathem. Publisher, v.41, 2002 (in Ukrainian)
- [24] Prykarpatsky Y.A., Samoilenko A.M., Prykarpatsky A.K. and Samoylenko V.Hr. *The Delsarte-Darboux type binary transformations and their differential-geometric and operator structure*. arXiv: math-ph/0403055 v 1 29 Mar 2004
- [25] Nimmo J.C.C. *Darboux transformations from reductions of the KP-hierarchy*. Preprint of the Dept. of Mathem. at the University of Glasgow, November 8, 2002, 11 p.
- [26] Prykarpatsky A.K. and Mykytiuk I.V. *Algebraic integrability of nonlinear dynamical systems on manifolds: classical and quantum aspects*. Kluwer Acad. Publishers, the Netherlands, 1998
- [27] Godbillon C. *Geometrie differentielle et mecanique analytique*. Paris, Hermann, 1969.
- [28] Teleman R. *Elemente de topologie si varietati diferentiabile*. Bucuresti Publ., Romania, 1964
- [29] De Rham G. *Varietes differentielles*. Hermann, Paris, 1955.
- [30] De Rham G. *Sur la theorie des formes differentielles harmoniques*. Ann. Univ. Grenoble, 1946, 22, p. 135-152.
- [31] Warner F. *Foundations of differential manifolds and Lie groups*. Academic Press, NY, 1971
- [32] Danford N. and Schwartz J.T. *Linear operators, v.2*. InterSci. Publ., NY, 1963
- [33] Datta B.N. and Sarkissian D.R. *Feedback control in distributed parameter gyroscopic systems: a solution of the partial eigenvalue assignment problem*. Mechanical Systems and Signal Processing, 2002, v.16, N.1, p.3-17
- [34] Gelfand I.M. and Shilov G.E. *Generalized functions and actions upon them*. Second edition. Moscow, Nauka Publisher, 1959 (in Russian)
- [35] Novikov S.P. (Editor) *Theory of solitons*. Nauka Publ., Moscow, 1980 (in Russian)
- [36] Pochynaiko M.D. and Sydorenko Yu.M. *Integrating some (2+1)-dimensional integrable systems by methods of inverse scattering problem and binary Darboux transformations*. Matematychni Studii, 2003, N20, p. 119-132

- [37] Zakharov V.E. and Shabat A.B. *A scheme of integration of nonlinear equations of mathematical physics via the inverse scattering problem. Part 1.* Func. Anal. and it Appl., 1974, v. 8,N3, p. 43-53; Part2. 1979, v.13,N3, p. 13-32 (in Russian)
- [38] Konopelchenko B.G. *On the integrable equations and degenerate dispersiopn laws in multidimensional soaces.* J. Phys. A: Math. and Gen., 1983, v. 16, p. L311-L316
- [39] Zakharov V.E. *Integrable systems in multidimensional spaces.* Lect. Notes in Phys., 1982, v.153, p. 190-216
- [40] Zakharov V.E. and Manakov S.V. *On a generalization of the inverse scattering problem.* Theoret. Mathem. Physics, 1976, v. 27, N3, p. 283-287
- [41] Levi D., Pilloni L. and Santini P.M. *Backlund transformations for nonlinear evolution equations in (2+1)-dimensions.* Phys. Lett, 1981, v.81A, N8, p. 419-423
- [42] Liu Wen. *Darboux transformations for a Lax integrable systems in 2n-dimensions.* arXiv:solve-int/9605002 v1 15 may 1996
- [43] Gu C.H. *Generalized self-dual Yahg-Mills flows, explicit solutions and reductions.* Acta Applicandae Mathem., 1995, v. 39, p. 349-360

\*)THE AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, DEPARTMENT OF APPLIED MATHEMATICS, KRAKOW 30059 POLAND, AND BROOKHAVEN NAT. LAB., CDIC, UPTON, NY, 11973 USA

*E-mail address:* yarchyk@imath.kiev.ua, yarpry@bnl.gov

\*\*)THE INSTITUTE OF MATHEMATICS, NAS, KYIV 01601, UKRAINE

\*\*\*)THE AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, DEPARTMENT OF APPLIED MATHEMATICS, KRAKOW 30059 POLAND

*E-mail address:* pryk.anat@ua.fm, prykanat@cybergal.com